



TERRY N. LONNER  
WILDLIFE BIOLOGIST



OF THE MONTANA CHAPTER

*of the* WILDLIFE SOCIETY



MONDAY AGENDA  
April 8, 1991

Ballroom B  
Chair: Marion Cherry

- 1:00        Opening Remarks
- 1:10        Habitat fragmentation and nest predation: emerging parallels between forest birds and waterfowl. Joe Ball. Montana Cooperative Wildlife Research Unit, University of Montana, Missoula, MT 59812.
- 1:40        Nesting biology of American avocets at two wildlife management areas in north central Montana. Kristi Dubois. U.S. Fish and Wildlife Service, Helena, MT 59626.
- 2:00        Double-crested cormorants and great blue herons: Management for biodiversity. Marcy Bishop. USF&WS, National Bison Range, Moiese, MT 59824.
- 2:20        Release techniques for Columbian sharp-tailed grouse in the Tobacco Valley, Montana. Lewis Young, David Genter, Bernie Hall, and Marilyn Wood. Kootenai National Forest, Eureka, MT 59717 (LY), The Nature Conservancy, Helena, MT 59624 (DG and BH), Montana Department of Fish, Wildlife and Parks, Kalispell, MT 59901 (MW).
- 2:40        Northern pocket gopher use of a black-tailed prairie dog colony following twelve years of no cattle grazing. Craig J. Knowles and Pamela R. Knowles. Department of Zoology, University of Montana, Missoula, MT and Fauna West Wildlife Consultants, Boulder, MT 59632.
- 3:00        Break
- 3:20        The blackfooted ferret (Mustela nigripes) reintroduction program with special reference to south Phillips County, Montana. Ron Crete and Ron Stoneberg, U.S. Fish and Wildlife Service, Helena, MT 59624 (RC), and Montana Department of Fish, Wildlife and Parks, Hinsdale, MT 59241 (RS).
- 3:40        Status of the swift fox in Montana. Craig J. Knowles and Pamela R. Knowles. FaunaWest Wildlife Consultants, Boulder, MT 59632.
- 4:00        Recent status of the bald eagle (Haliaeetus leucocephalus) in Montana. Dennis L. Flath and Robert M. Hazelwood. Montana Department of Fish, Wildlife and Parks, Bozeman, MT 59717 (DF) and U.S. Fish and Wildlife Service, Helena MT 59626 (RMH).
- 4:20        Status, distribution, and ecology of the golden eagle in Yellowstone National Park. Terry McEneaney, Yellowstone National Park, WY 82190.



MONDAY AGENDA  
April 8, 1991

Ballroom C  
Chair: Dick Hutto

- 1:00           Opening Remarks
- 1:10           An examination of bird distribution in old-growth and rotation-aged ponderosa pin/Douglas-fir stands from a landscape perspective. Sallie J. Hejl, Jock S. Young, Susan A. Colt, and Ruth E. Woods. USFS, Intermountain Experiment Station, Missoula, MT 59801.
- 1:40           On the conservation of neotropical migratory land birds in Montana. Richard L. Hutto. Division of Biological Sciences, University of Montana, Missoula, MT 59812.
- 2:00           Foraging site selection by insectivorous forest birds in western Montana. Andrew H. Bosma. Division of Biological Sciences, University of Montana, Missoula, MT 59812.
- 2:20           Snag use for nest site selection of four primary cavity nesting woodpeckers in the Bridger Mountains, Montana. James R. Sparks. Biology Department, Montana State University, Bozeman, MT 59715, and Gallatin National Forest, Big Timber, MT 59011.
- 2:40           Logging and forest birds: community-level patterns and a critical measure of habitat quality. Bret W. Tobalske. Division of Biological Sciences, University of Montana, Missoula, MT 59812.
- 3:00           Break
- 3:20           Small mammal density and diversity in Douglas-fir (Pseudotsuga menziesii) old-growth forests as related to forest structure. Patricia Cramer. Biology Department, Montana State University, Bozeman, MT 59717.
- 3:40           Red squirrel (Tamiasciurus hudsonicus) cache sites and cache pattern in Pattee Canyon, Montana. Rebecca S. Burton. Division of Biological Sciences, University of Montana, Missoula, MT 59812.
- 4:00           Rediscovery of the spotted bat (Euderma maculatum), pallid bat (Antrozous pallidus) and Townsend's big-eared bat (Plecotus townsendii) in south central Montana. David Worthington. Division of Biological Sciences, University of Montana, Missoula, MT 59812.
- 4:20           Results of harlequin duck surveys on the Flathead National Forest, Montana. John C. Carlson. Montana Natural Heritage Program, Helena, MT 59620 and University of Wyoming, Laramie, WY.
- 4:30           Poster Session



TUESDAY AGENDA  
April 9, 1991

Ballroom B and C

8:00 Opening remarks

8:15 Panel discussion "Are we managing for all wildlife"  
Moderator: Dr. Lee Metzgar

10:00 Break

Chair: Fritz Prellwitz

10:20 Dynamics of beaver food caches and cache size as a predictor of colony size in Wyoming. Cynthia Osmundson and Steven Buskirk. U.S. Fish and Wildlife Service, CMR National Wildlife Refuge, Box 110, Lewistown, MT 59457 (CO), Department of Zoology and Physiology, University of Wyoming, Laramie, WY 82071.

10:40 Mortality rates of the pine marten in southwest Montana. Craig Fager. Department of Biology, Montana State University, Bozeman, MT 59717.

11:00 Wolf ecology update. Dan Pletcher. School of Forestry, University of Montana, Missoula, MT 59812.

11:20 Mortality and seasonal distribution of white-tailed deer in an area recently recolonized by wolves. Jon S. Rachael, School of Forestry, University of Montana, Missoula, MT 59812.

11:40 The significance of disease in wildlife populations. Mark Johnson DVM. Wildlife Veterinary Consultant, Gardiner, MT. 59030.

12:00 Lunch Break

Chair: Mike Aderhold

1:00 Survival rates of black bear in northwest Montana. Wayne F. Kasworm and Timothy J. Thier. U.S. Fish and Wildlife Service, 475 Fish Hatchery Road, Libby, MT 59923.

1:20 Summary of questionnaire given to spring black bear hunters in northwestern Montana. Timothy J. Thier. U.S. Fish and Wildlife Service, 475 Fish Hatchery Road, Libby, MT 59923.

1:40 Grizzly bear population augmentation in the cabinet mountains, northwest Montana. Wayne F. Kasworm, Timothy J. Thier, and Christopher Servheen. U.S. Fish and Wildlife Service, Libby, MT 59923 (WK and TT), U.S. Fish and Wildlife Service, University of Montana, Missoula MT 59812 (CS).



TUESDAY AGENDA (Cont.)

- 2:00 Increasing mountain lion populations and human-lion interactions in Montana Keith E. Aune. Montana Department of Fish, Wildlife and Parks, Mont. State Univ. Bozeman, MT 59717.
- 2:20 Moose-habitat relationships in northwestern Montana and southeastern British Columbia. Margaret Langley, School of Forestry, University of Montana, Missoula, MT 59812.
- 2:40 Some population characteristics of the Yellowstone bison herd. Helga Ihsle Pac and Kevin Frey. Montana Department of Fish, Wildlife and Parks, Mont. State Univ. Bozeman, MT 59717.
- 3:00 Break

Chair: Keith Aune

- 3:20 Summer habitat use of white-tailed deer in northwestern Montana. John Morgan. Department of Biology, Montana State University, Bozeman, MT 59717.
- 3:40 The Flying D Ranch and its significance to wildlife management in southwest Montana. Kurt Alt. Montana Department of Fish, Wildlife and Parks, Bozeman, MT 59715.
- 4:00 Home range use of historical and present elk populations in south central Montana. Fred G. VanDyke and John P. Skubinna. Montana Department of Fish, Wildlife and Parks, Red Lodge, MT 59068.
- 4:20 Elk plus cows times politics equals ? Kurt Alt, Fred King, Mark Petroni, and Ron Schott. Montana Department of Fish, Wildlife and Parks, Bozeman, MT 59715 (KA and FK), U. S. Forest Service, Beaverhead National Forest, Ennis, MT 59729.

TUESDAY EVENING

Montana Chapter Business Meeting	6:00-7:00	Gran Tree Ballroom
Western Barbecue Buffet and Social Hour	7:00-8:00	Gran Tree Atrium
Awards Presentation	7:30-8:00	
Entertainment	8:00-9:00	
(Charlie Russell Yarns)		



HABITAT FRAGMENTATION AND NEST PREDATION: EMERGING PARALLELS BETWEEN FOREST BIRDS AND WATERFOWL

BALL, I. J., Montana Cooperative Wildlife Research Unit, U.S. Fish and Wildlife Service, University of Montana, Missoula, MT 59812

Abstract

Low nest success and faltering populations are a serious threat to neotropical migrant songbirds in the eastern United States and upland nesting ducks in the Prairie Pothole Region. In both cases, excessive nest predation occurs because of habitat fragmentation and supplementation of predator populations by humans. Viable recruitment rates among both groups seem to occur only in the very largest remaining parcels of unfragmented habitat. Protecting or providing "good" habitat in relatively small parcels does not solve, and may exacerbate, the problem. Acknowledging this problem and developing effective and socially acceptable solutions will be difficult, but doing so offers important common ground to those interested in biological diversity, nongame wildlife, and game species.

NESTING BIOLOGY OF AMERICAN AVOCETS AT TWO WILDLIFE MANAGEMENT AREAS IN NORTHCENTRAL MONTANA.

Kristi DuBois, U.S. Fish and Wildlife Service, Fish and Wildlife Enhancement, 301 S. Park, Drawer 10023, Helena, MT 59626

Abstract

American Avocet populations were studied at Benton Lake National Wildlife Refuge and Freezout Lake State Wildlife Management Area during the 1990 breeding season, as part of a study to assess the potential impacts of selenium from irrigation drainwater on water bird reproduction. A total of 103 nests were monitored at Benton Lake and 185 nests were monitored at Freezout Lake. Nest success within different avocet colonies varied from 25% to 100%. Overall nest success was 69% at Benton Lake and 64% at Freezout Lake. Mammalian predation and flooding were the primary causes of nest loss. Average clutch size was 3.8 eggs/clutch for both areas. Nesting chronology, egg size, and young size were compared between the two areas. Survival of young to fledging appeared to be good at both areas.



DOUBLE-CRESTED CORMORANTS AND GREAT BLUE HERONS - MANAGEMENT FOR BIODIVERSITY

Marcy Bishop, USFWS, National Bison Range, Moiese, MT 59824

Abstract

Wetland management at Ninepipe and Pablo National Wildlife Refuges has provided for biodiversity and for long term research on a cormorant and heron colony under undisturbed conditions. Management includes protection from outside intrusion, timing of other management duties, restoration and protection of nesting islands and trees, passive predator control and careful banding and study methods. This program has provided for 20 years of data on nesting success, population dynamics, nest stratification, resource partitioning, interactions with other species, behavior and fledging of young as they occur in a natural, undisturbed colony.

RELEASE TECHNIQUES FOR COLUMBIAN SHARP-TAILED GROUSE IN THE TOBACCO VALLEY, MONTANA

Lewis Young, Kootenai National Forest, Eureka, MT 59917  
David Genter, Montana Natural Heritage Program, Helena, MT 59620  
Bernie Hall, The Nature Conservancy, Helena, MT 59620  
Marilyn Wood, Montana Department of Fish, Wildlife, and Parks, Kalispell, MT 59901

*Abstract*

Since 1987, 61 Columbian sharp-tailed grouse have been transplanted from British Columbia, Canada, to the Tobacco Valley in northwestern Montana to sustain and rejuvenate the last known lek in northwestern Montana. Each year, 10-20 grouse were trapped on leks in Canada using both drop nets and walk-in traps, then transported by small aircraft or ground transportation as soon as possible. Birds were released on the leks in the presence of other grouse using custom made release boxes while sharp-tail vocalizations were played on an audio tape player. Releases were made both morning and evening. Observations included leg bands and in 1990, radio-collars, indicate the transplanted birds join the leks and have good survival rates.



Craig J. Knowles and Pamela R. Knowles  
Dept. of Zoology, University of Montana,  
Missoula, MT 59812 (CJK)  
FaunaWest Wildlife Consultants, P.O. Box 113,  
Boulder, MT 59632 (PRK)

Abstract: Northern pocket gophers (Thomomys talpoides) were found to coexist with black-tailed prairie dogs (Cynomys ludovicianus) within a portion of a prairie dog colony excluded from livestock grazing for a period of twelve years. Relative density of fresh pocket gopher mounds inside the enclosure was 60.8/km while relative density outside the enclosure was 4.5/km. Forbs dominated the vegetation in the area used by pocket gophers. Forb and grass production were greater within the enclosure than the adjacent area outside the enclosure.

The black-footed ferret (Mustela nigripes) reintroduction program with special reference to south Phillips County, Montana

Ron Crete, USFWS, P.O. Box 10023, Helena, MT 59626

Ron Stoneberg, MDFWP, Box 424, Hinsdale, MT 59241

Abstract: The black-footed ferret (BFF) is one of the rarest and most endangered mammals in North America. It is listed as endangered under Montana and national endangered species laws. All known members of the species are in captivity. An aggressive captive breeding program is reaching the goal of producing individuals in excess of the captive breeding objectives for reintroduction into the wild planned for 1991. National and state interdisciplinary teams are seeking recovery objectives of the BFF Recovery Plan. Surveys to find potential reintroduction sites throughout the historical range of the species identified South Phillips County as a candidate for ferret recovery actions. Planning for BFF recovery in northcentral Montana is dove-tailed with Bureau of Land Management resource planning. An interagency team of biologists works with individuals and groups in this Montana potential recovery area and statewide to understand concerns about black-footed ferret recovery. They will draft a BFF reintroduction and management plan in Spring 1991 aimed to avoid or mitigate impacts to the local economy and desired lifestyles of Montanans while meeting state and national responsibilities for the conservation of the black-footed ferret.



## Status of the Swift Fox in Montana

Craig J. Knowles and Pamela R. Knowles  
FaunaWest Wildlife Consultants, P.O. Box 113,  
Boulder, MT 59632

Abstract: A search was made for historical and recent records of the swift fox (Vulpes velox) in Montana. The first observations of swift foxes in Montana were made by the Lewis and Clark expedition in 1805 along the Marias River and at the Great Falls of the Missouri River. Forty-four specimens were collected prior to 1910. Forty-three of these were collected at Kipp and Blackfoot in upper Teton County and one was collected at Fort Benton. The last historic observations of the swift fox in Montana were made by Bailey and Bailey in 1918 on the plains along the eastern edge of Glacier National Park. The lack of any confirmed records since 1918 prompted Hoffmann et al. in 1969 to declare the swift fox extinct in Montana. However, beginning in 1978, scattered reports of swift foxes in Montana have been documented. Three records have come from southeastern Montana near Miles City, Glendive, and Broadus. Three unconfirmed sightings have been reported near Circle, Forsyth, and Miles City. In addition, swift foxes reintroduced in Alberta and Saskatchewan, Canada have been documented to cross into Montana north of Havre. Two established swift fox populations exist near Montana, and are located near Gillette, Wyoming and in Perkins County in northwestern South Dakota.

### RECENT STATUS OF THE BALD EAGLE (Haliaeetus leucocephalus) IN MONTANA

Dennis L. Flath, Montana Department of Fish, Wildlife and Parks,  
FWP Building, MSU Campus, Bozeman, MT 59717;

and,

Robert M. Hazlewood, US Fish and Wildlife Service, Federal Building  
and US Courthouse, 301 South Park, PO Box 10023, Helena, MT 59626

### Abstract

Changes in the status of Montana's bald eagle nesting population are documented, including factors responsible for population depression and evident recovery. Productivity, rate of population increase and attributes of nesting habitat are described. The nesting population increased from 26 territories producing 29 young in 1980 to 108 territories producing 130 young in 1990. The first North American record of bald eagles producing 4 young in one brood is documented. Management problems and conservation recommendations are presented.



# Status, Distribution, and Ecology of the Golden Eagle in Yellowstone National Park

Terry McEneaney, Yellowstone National Park,  
P.O. Box 168, Yellowstone Nat. Park, Wyo. 82190

## Abstract

The golden eagle (*Aquila chrysaetos*) is one of the most efficient avian predators in the western United States, yet its status in Yellowstone has remained relatively unknown until now. Incidental golden eagle data was collected while censusing for peregrine falcons in Yellowstone from 1986 - 1990. Remote areas of Yellowstone were initially surveyed by helicopter and checked for greater detail using ground reconnaissance. An update on the status, distribution, and ecology of the golden eagle will be discussed along with a historical perspective of golden eagle activity in Yellowstone.

An Examination of Bird Distribution in Old-growth and Rotation-aged Ponderosa Pine/Douglas-fir Stands from a Landscape Perspective

Hejl, Sallie J., Jock S. Young, Susan A. Colt, and Ruth E. Woods, U.S.F.S.,  
Intermountain Research Station, Missoula, Montana

In each of 16 old-growth (200+ years) and 16 rotation-aged (80-120 years) ponderosa pine/Douglas-fir stands in western Montana and adjacent Idaho, birds were counted four times during the 1989 breeding season. Of the 68 species recorded, 14 were found exclusively in old-growth and seven exclusively in rotation-aged stands. For the 26 common species, (1) 8 species were more abundant in old-growth, (2) 7 species were more abundant in rotation age, and (3) 11 were not associated significantly with either habitat. In order to examine landscape-level associations, we analyzed the distribution of the 26 common species in relation to landscape variables calculated from a 2 square-mile area centered on each stand. Landscape variables, especially total amount of old-growth forest and amount of open land, helped to explain the distribution of 14 of the common species.

g  
6

n  
n  
e  
g  
t  
d



## ON THE CONSERVATION OF NEOTROPICAL MIGRATORY LANDBIRDS IN MONTANA

Richard L. Hutto, Biological Sciences, University of Montana, Missoula, MT 59812

### Abstract

I will describe the neotropical migratory landbird conservation initiative developed by federal and state agencies and non-governmental organizations in December 1990. Montana is in an excellent position to accomplish a number of the goals associated with this program, especially with respect to the development of monitoring methods, and the use of existing GIS databases to build habitat suitability models that incorporate landscape-level information. In short, neotropical migrants have the potential to serve as a powerful "management indicator group".

## SNAG USE FOR NEST SITE SELECTION OF FOUR PRIMARY CAVITY NESTING WOODPECKERS IN THE BRIDGER MOUNTAINS, MONTANA

Sparks, James, R., Biology Department, Montana State University, Bozeman, Montana 59715 / Big Timber Ranger District, Gallatin National Forest, Big Timber, Montana 59011.

### Abstract

During June and July of 1988, 1989, and 1990, 46 active nests of 4 primary cavity nesting woodpeckers were located in Douglas-fir (*Pseudotsuga menziesii*) forests in the Bridger Mountains of southwestern Montana. 12 Williamson's Sapsucker, 10 Hairy Woodpecker, 9 Three-toed Woodpecker, and 15 Northern Flicker nests were discovered. 31 Douglas-fir, 8 lodgepole pine (*Pinus contorta*), 4 subalpine fir (*Abies lasiocarpa*), 2 limber pine (*Pinus flexilis*), and 1 Engelmann spruce (*Picea engelmannii*) were selected as cavity nesting trees. All trees utilized for nest sites were standing dead trees (snags). To minimize interspecific competition for cavity trees during the breeding season, these 4 woodpecker species appear to select different tree species, select different size (diameter and height) snags, or select snags in different stages of decay for nest sites.



# THE STATUS OF THE HARLEQUIN DUCK ON 5 NATIONAL FORESTS OF WESTERN MONTANA

John C. Carlson, Montana Natural Heritage Program, Helena, MT 59620, and University of Wyoming, Laramie, WY 82071

Status surveys for harlequin ducks were conducted on portions of 5 National Forests in Montana during the 1990 breeding season. A total of 91 streams were surveyed on the Lewis & Clark, Flathead, Gallatin, Kootenai, and Lolo National Forests. Stream flow and habitat characteristics were obtained on streams surveyed on the Gallatin, Flathead, Kootenai, and Lolo National Forests. A minimum of 101 individual birds were located on 20 streams and rivers; 25 were males, 25 females, and 51 young of the year. The average brood size was 3.18 young. Harlequin ducks were found predominantly on low gradient, braided type streams with stretches or side channels of slower water. Additional sightings on the Rocky Mountain Division of the Lewis & Clark National Forest revealed an estimated 80 - 122 harlequin ducks, with 13 - 22 broods observed last summer.

## DYNAMICS OF BEAVER FOOD CACHES AND CACHE SIZE AS A PREDICTOR OF COLONY SIZE IN WYOMING

CYNTHIA OSMUNDSON, U.S. Fish and Wildlife Service, Charles M. Russell National Wildlife Refuge, Box 110, Lewistown, MT 59457  
STEVEN BUSKIRK, Department of Zoology and Physiology, University of Wyoming, Laramie, WY 82071

Abstract: This study investigated the relationship between the size of food caches and the number of beaver in a colony. Two recent studies suggested that cache size may indicate colony size. Results of our study did not support this reported relationship; the correlation between cache size and colony size was not significant ( $r = 0.16$ ,  $P = 0.24$ ). Cache construction began in late September and caches grew at a mean rate of  $0.45 \text{ m}^3/\text{week}$  until final freeze-up, typically during the first week of November. Mean final cache area was  $61 \text{ m}^2$ . Colony size was estimated during night observations with infra-red night vision devices. Mean colony size was estimated to be 5.05. Late fall movements and cache abandonment were identified and studied through the use of implanted radio transmitters. The ability of observers to locate and accurately estimate size of caches from the air was assessed. Aerial observers correctly classified caches by size 47.5% of the time, whereas 11% of known caches were overlooked. Estimated availability of cache material was not significantly correlated with cache size ( $r = 0.02$ ,  $P = 0.93$ ).



MORTALITY RATES OF THE PINE MARTEN IN SOUTHWEST MONTANA  
Craig Fager, Montana State University  
Quinton Kujula, Montana State University

Abstract: Trapping mortality in pine marten was studied during the 1989-1990 and 1990-1991 trapping seasons through mark-recapture methods in the upper Big Hole and West Yellowstone areas of southwest Montana. Trappers caught 40% (2 of 5) of radio collared marten in the upper Big Hole study area during the 1989-1990 season. The marked sample represented 12.5% of the known study area harvest (n=16). This area was abandoned by trappers during the 1990-1991 trapping season. Eight of 23 marked marten were captured by trappers in the West Yellowstone study areas. The marked sample represented 28.6% of known study area harvest (n=28). Harvest sex ratios varied from 2.2M:1F in the upper Big Hole to 1.54M:1F in the West Yellowstone areas. These ratios differed from pre-season live trapping sex ratios (1M:1F, 1M:1.3F) but the difference harvested was not statistically significant. The radio collared marten harvested in the upper Big Hole were trapped approximately 5 linear kilometers from their point of live capture. Radio collared marten harvested in the West Yellowstone area were all captured within 0.5 linear kilometers of their point of live capture during the 1989-1990 season and 0.6-5.4 km during the 1990-1991 season. Variations in harvesting techniques between study areas may explain some of the differences in sex ratios and movements of harvest marten. Trappers in the West Yellowstone area employ a much greater trap density per unit area (approximately 6 sets/km road or trail) than the individual in the upper Big Hole (approximately 1 set/km road or trail).



## INTRODUCTION

The Southwest Montana Pine Marten Project is a Montana Department of Fish, Wildlife, and Parks and U.S. Forest Service supported project designed to characterize pine marten habitat and harvest throughout southwest Montana. Recent concerns over an apparent decrease in the harvestable surplus of marten populations were a driving force behind the initiation of the investigation. The project is scheduled to run 3 years. Because marten are easily harvested but are believed to live in extremely specialized habitats, habitat and harvest questions cannot be easily separated; however, for the purposes of this paper, the 2 questions are assumed to be largely independent. This paper combines information gathered by 2 separate investigators during the first 2 field seasons of the Southwest Montana Pine Marten Project.

## STUDY AREAS AND METHODS

Three study areas were used to evaluate marten harvest levels. Historical harvest levels, access, and habitat manipulations were all considered in study area selection.

The 32 km<sup>2</sup> Beaver Creek study area lies in hunting district 362, northwest of Quake Lake. Vehicle access is limited to less than 10 km, but 4 trailheads into the Taylor Hilgard Wilderness and adjacent roadless areas provide access to a vast area. Timber harvesting activity has been severely restricted in this area. Historically, 17 to 20 marten have been taken annually from the drainage by a single trapper.

The 68 km<sup>2</sup> West Yellowstone Flats study area lies immediately north of West Yellowstone in hunting district 361. The area is bound by Yellowstone National Park on the eastern edge, Cougar Creek on the north, and Hebgen Lake on the western edge. The flat, open terrain make vehicle access almost unrestricted, particularly with snow machines. Trapper numbers as well as harvest are both highly variable in this area. The known 1989-1990 harvest was 1 marten, but personal averages per trapper in the 7 to 10 range appear to be more normal. This study area was the only one to be directly affected by fire during the summer of 1988.

The 153 km<sup>2</sup> upper Big Hole study area lies northwest of the Big Hole Battlefield, in hunting district 321, and includes the Bender, Schultz, Johnson, and Tie Creek drainages. The area is bound by the Bitterroot National Forest to the northwest and Highway 43 on its southern edge. Road building and timber harvesting activity have been heavy in the last several decades. Several U.S.F.S. trails found in the area are also utilized by trappers. Historically, this area has been regarded as some of southwest Montana's best marten producing habitat. Individual trappers have harvested nearly 50 marten from the study area in some years. During the 1980's, the study area has been trapped intermittently by



at least 3 different individual during different years.

Marten were live-trapped prior to the opening of the December 1 trapping season with single door, wire mesh traps. Through December 1991, most marten were fitted with 148MHZ AVM type P2-B transmitters and eartags. Marten captured in remote, inaccessible areas were fitted with eartags only. Limited live trapping was conducted during the trapping season on the West Yellowstone Flats and upper Big Hole study areas. Because areas live trapped and areas harvested did not completely overlap in all cases, each marked marten was evaluated with regard to harvest vulnerability. Trappers were contacted on an individual basis both before and during the trapping season. These individuals provided valuable information concerning trap density, techniques, and capture locations for both marked and unmarked marten.

Baseline population productivity and density indexes for local populations were developed through 1 km track transects, live trapping-harvest success rates, and home range analysis. Differences in trapping methods other than trap density were not considered as variables.

## Results and Discussion

Population indices for the 1989-1990 field season suggest that marten populations in all three study areas were severely depressed. Relative track densities from permanent track transects all fell below 0.6 tracks/km, and were consistent with densities found in an Ontario population suffering from a food shortage (Thompson and Colgan 1987). Harvest data from the trappers utilizing the study areas provided additional evidence of population depression. Success rates in the 1988-1989 season were reduced by 70-90% compared to the 1989-1990 season. (Whitman pers. comm.; Depas pers. comm.). Live trapping success and home ranges would also seem to indicate depressed populations when compared with other studies. Approximately 2.8 times the trapping effort was required to capture marten in our study areas than was required by Hawley and Newby (1957) in another Montana study. Inflated home ranges may be a sign of additional stress on marten populations in Southwest Montana. Home ranges calculated to date run (Table 1) 3 to 10 times as large as home ranges reported for Alaskan marten (male average 6.62 km<sup>2</sup>; female average 3.71 km<sup>2</sup>) reported by Burskirk (1983).



Table 1. Ninety-five percent, 75%, and core areas<sup>a</sup> for 9 marten, by sex and season, from the upper Big Hole and West Yellowstone study areas.

I.D.	Sex	Season	n	Home range in km <sup>2</sup>		
				95%	75%	Core
West Yellowstone						
521	M	Winter	28	25.7	9.04	5.6
205	M	Win./Sp.	15	68.0	28.4	13.25
727	F	Winter	8	4.9	3.37	
845	M	Winter	24	29.2	17.7	8.6
845	M	Spring	31	56.6	24.7	17.71
845	M	Win./Sp.	55	67.0	31.36	18.3
Upper Big Hole						
970	F	Winter	29	7.2	3.0	1.8
762	M	Win./Sp.	10	19.1	12.78	3.8
492	M	Winter	18	9.7	5.5	3.5


<sup>a</sup> Core areas reflect approximately 50% utilization.

Figure 1 shows harvest and price trends from the 1979-1980 through 1989-1990 trapping seasons. An average of 428 marten were harvested over the 11 year period. In response to concerns over reduced harvest in some of Region 3's best marten producing habitats, the November through February season was reduced in 1987-1988 to include only the months of December and January. The 1990-1991 harvest (157 marten) appears significantly lower than any of the preceding 11 years. Price data were not available and no marten were bought in 1990-1991 by the principal fur buyer in the Bozeman area. The number of trappers participating annually appears to be declining across the region; however, not enough data are available at this time to draw any conclusions.

Table 2 shows harvest by study area. Harvest rates in 1989-1991 were from 70 to 90% lower than harvests taken on the same areas earlier in the 1980's. Part of this may be because of the 50% reduction in season length. However, given the similar efforts put forth between seasons, it is unlikely that this explains more than a small percentage of the seasonal differences. Losing the month of November has been cited by several trappers we contacted as the primary reason for lower catches during high marten years, after the season length was reduced.



Figure 1. Marten Harvest and Prices  
1979-1990



Data from Howard Hersh, MDFWP

Table 2. Harvest statistics by study area.

Study area	Harvest	No. Marked In Harvest	% of Harvest	Historic Level
Beaver Creek (H.D. 362)				
1989-1990	6	3 of 11	50%	17-20
1990-1991	16	3-4 of 6	19-25%	
West Yellowstone Flats (H.D. 361)				
1989-1990	1	1 of 1	100%	7-10/Indiv.
1990-1991	6	1 of 6	17%	
Upper Big Hole (H.D. 321)				
1989-1990	16	2 of 5	12.5%	30-50
1990-1991		Abandoned by trapper		

Marked marten constituted between 12.5% and 100% of the known harvest by study area. These values do not represent true harvest percentages because areas live trapped and areas harvested did not overlap 100% in the study areas of the West Yellowstone Flats and the upper Big Hole. However, the values do illustrate that not all marten are likely to be killed by trappers operating in their ranges despite the susceptibility of marten to trapping because all marked marten were at least exposed to trap lines as judged through location of live



capture during November as well as radio locations during the trapping season.

Marten possess a very limited ability to avoid traps in their immediate activity centers. Survivorship of marten that remained resident in the immediate vicinity of the point of live capture was further diminished if local trappers were employing high densities of traps. Marked marten in the Big Hole were harvested 5 km from their point of live capture under a trapping scheme with approximately 1 trap/km. Marten in the West Yellowstone area, under a much higher trap density of at least 6 traps/km, were with 1 exception, caught within 1.5 km of their point of live capture. Radio telemetry has provided largely inconclusive data on marten live trapped in November, but it is clear that a large portion of the Beaver Creek and Big Hole November samples avoided both further live capture and harvest through dispersal after initial live capture. Small sample sizes preclude any meaningful statistical analysis on these samples.

Pre-trapping season sex ratios from all study areas approached 1:1 (Big Hole=1M:1F, West Yellowstone=1M:1.3F). These values are different, but not statistically, from the harvest sex ratios of 2.2M:1F in the Big Hole and 1.54M:1F in the West Yellowstone area.

#### Literature Cited

- Buskirk, S.W. 1983. The ecology of marten in southcentral Alaska. Ph.D. Thesis, Univ. Alaska, Fairbanks. 131pp.
- Hawley, V.D. and F.E. Newby. 1957. Marten home ranges and population fluctuations. J. Mammal. 38(2):174-184.
- Soutiere, E.C. 1978. The effects of timber harvesting on marten. M.S. Thesis. Univ. Maine, Orono. 62pp.
- Thompson, I.D. and P.W. Colgan. 1987. Numerical responses of martens to a food shortage in northcentral Ontario. J. Wildl. Manage. 51(4):824-835.







RED SQUIRREL (Tamiasciurus hudsonicus) CACHE SITES AND CACHE  
PATTERN IN PATTEE CANYON, MONTANA

Burton, Rebecca S., Division of Biological Sciences,  
University of Montana, Missoula, MT 59812

ABSTRACT

The spatial pattern and physical attributes of seed caches used by red squirrels (Tamiasciurus hudsonicus) were the focus of this study conducted in Pattee Canyon, near Missoula, Montana. Within the 320 X 290-m study site there were 103 caches. Each of the squirrels studied used at least 10 caches. There was no evidence of cone theft.

In comparison with unused sites, cache sites were associated with soft soil and were near large ponderosa pine (Pinus ponderosa) trees. Caches had more vegetative cover around the center of the cache than at 3- or 5-m from the center. Unused sites had more total cover than used sites. Caches were not significantly closer to nests or caches than were unused sites and were not significantly associated with tree or log density.

RE-DISCOVERY OF THE SPOTTED BAT (Euderma maculatum), PALLID  
BAT (Antrozous pallidus), AND TOWNSEND'S BIG-EARED BAT (Plecotus  
townsendii) IN SOUTH-CENTRAL MONTANA

David Worthington, Division of Biological Sciences, University of  
Montana, Missoula MT 59812

Abstract

The spotted bat and pallid bat were known previously in Montana from one individual each captured in 1949 and 1978, respectively. The Townsend's big-eared bat is known to breed in Montana, and was known previously in south-central Montana from winter hibernacula. Two spotted bats, 26 pallid bats, and 11 Townsend's big-eared bats were observed in 1989 and 1990 in the Pryor Mountains. Juveniles and lactating females were observed in all three species, suggesting that the species breed in the region. All three species are listed as species of special concern by the Montana Natural Heritage Program, while the spotted bat and Townsend's big-eared bat are listed under category 2 as candidates for the Endangered Species Act and as sensitive species by the USFS.



LOGGING AND FOREST BIRDS: COMMUNITY-LEVEL PATTERNS AND A CRITICAL MEASURE  
OF HABITAT QUALITY

Tobalske, Bret W. Division of Biological Sciences, University of Montana,  
Missoula, MT 59812

Abstract

I studied the influence of logging upon a community of breeding birds in western larch/Douglas-fir stands on the Coram Experimental Forest, Montana. Ten species, four foraging guilds, and two nesting guilds showed significant differences in abundance among stand conditions during 1989 and 1990. Interactions for abundance between stand conditions and years were evident for two species, one foraging, and one nesting guild. In general, the patterns observed were intuitive. Conifer tree nesters, for example, were least abundant in clearcuts. Ironically, cavity nesters, including Red-naped Sapsuckers, were equally abundant in logged (with some trees and snags reserved) and unlogged habitat. I studied reproductive success among sapsuckers during 1990 to evaluate the suitability of disturbed habitat for this species, and observed higher productivity in logged stands. These data validate current guidelines regarding snag retention within cutting units, and highlight the importance of paper birch and aspen for cavity nesting and other tree-dependent bird species.

SMALL MAMMAL DENSITY AND DIVERSITY IN DOUGLAS FIR  
(PSEUDOTSUGA MENZISII) OLD GROWTH FORESTS AS  
RELATED TO FOREST STRUCTURE

Patricia Cramer  
Biology Department, Conservation Biology Program, Montana State  
University, Bozeman, MT 59717

Abstract

Density and diversity of small mammals in old growth forests were studied during the summer months of 1990. Live trapping methods were used on two similar Douglas fir old growth forests in the Hyalite Basin of the Gallatin Range, Montana. The two forest types were similar in aspect, slope, tree composition and age, but differed in complexity of structure, particularly with respect to the amount of downed material contributing to fuel loading. The more complex forest averaged 60.5 tons of downed logs per acre and after 3,600 trap-nights, 317 individual animals were caught there. The more simple forest averaged 6.9 tons of downed logs per acre and produced 192 individual animals after 3,600 trap-nights. Results of this study may indicate that dead and downed logs play a significant factor in numbers of small mammals in old growth forests. These small mammals serve as a prey base for other animals in old growth areas, in particular, the pine marten relies heavily on red-backed voles which constituted a majority of animals caught in this study.



## MORTALITY AND SEASONAL DISTRIBUTION OF WHITE-TAILED DEER IN AN AREA RECENTLY RECOLONIZED BY WOLVES

JON S. RACHAEL, School of Forestry, University of Montana,  
Missoula, MT 59812

**Abstract:** White-tailed deer (*Odocoileus virginianus*) are the most abundant prey item of wolves (*Canis lupus*) that have recently recolonized northwestern Montana, but little is known about mortality or seasonal movements of deer in this area. Since January 1990, I have been investigating cause-specific mortality, seasonal distribution, and fawning-site selection of white-tailed deer in this region. Of 38 female white-tailed deer captured and radio-collared during winter 1989-1990 and 1990-1991, 2 were killed by wolves. A bear (*Ursus americanus*), a coyote (*C. latrans*), and a mountain lion (*Felis concolor*) each killed 1 radio-collared deer, and humans killed 2. Predators killed 3 radio-collared deer during winter and 2 during the fawning period in early June, but killed none during summer or autumn. Although 5 radio-collared deer spent the summer within 1 km of a wolf den in 1990, wolves killed none, possibly because fawns typically comprise a large portion of the diet of wolves during this period. Deer captured on 3 winter ranges in western Glacier National Park migrated in all directions an average of 12 km ( $n = 21$ , Range = 0 to 40 km) to summer ranges. Field work will continue through summer 1991.

Although once common throughout the western United States, the gray wolf was extirpated from the Northern Rocky Mountains by widespread public and private control efforts. Reported sightings of wolves were extremely rare in Montana by the 1930's and remained sporadic through the late 1970's (Day 1981). Occasional wolf sightings were probably of dispersing or lone wolves. No documented cases of wolf reproduction in the western U.S. occurred until 1986 when a den was found in Glacier National Park (Ream et al. 1987, 1989). Another den was found in Glacier Park in 1987, and 2 dens were located in British Columbia within 10 km of the international border in 1988. Wolves denned within Glacier National Park again in 1989, but the litter failed. In 1990 wolves denned at 2 sites in Glacier National Park and produced 12 pups. By December 1990, 34 wolves were known to inhabit Glacier National Park and the immediate surrounding areas of the Flathead National Forest and southeastern British Columbia.

The on-going natural recolonization of wolves in northwestern Montana has occurred in an area unlike many other areas where wolf studies have been conducted. Because this area has been without a breeding population of wolves for more than 50 years, ungulate populations in the relative security of Glacier National Park have probably reached an equilibrium with their habitat and other predators. Additionally, prey diversity in this system is higher than in most other systems studied. Most studies of wolf-prey interactions elsewhere have been conducted in areas with only 1 or 2 primary prey species (e.g. Murie 1944,



Mech 1966, Messier and Crete 1985, Ballard et al. 1987). White-tailed deer, mule deer (*O. hemionus*), elk (*Cervus elaphus*), and moose (*Alces alces*), are relatively abundant and provide a potential prey base for wolves in northwestern Montana. Low numbers of mountain goats (*Oreamnos americanus*) and bighorn sheep (*Ovis canadensis*) are present at higher elevations.

Wolf recolonization is highly controversial. Researchers studying public attitudes toward wolves have documented a public concern for native ungulate populations (e.g. Kellert 1985, McNaught 1987, Bath 1987, Bath and Buchanan 1989, Tucker and Pletscher 1989). To answer questions from the public and to justify management actions, resource managers need reliable information on impacts of wolf predation on ungulate populations.

In the North Fork drainage of the Flathead river in northwestern Montana, nearly half (45%) of the prey killed by wolves were white-tailed deer (Boyd et al. in prep.). Although white-tailed deer are the top management priority of the Montana Department of Fish, Wildlife and Parks in northwestern Montana, cause-specific mortality rates and seasonal movement patterns of deer in this area were unknown.

Results of a predator-prey study in an area being recolonized by wolves may yield valuable information with applications to other areas where wolves may recolonize or be reintroduced. My research objectives were to:

- 1) Evaluate cause-specific mortality of white-tailed deer within the area recolonized by wolves;
  - 2) Document seasonal distribution of white-tailed deer, including identification of key areas of seasonal use;
  - 3) Initiate an index to monitor deer abundance over time;
- and,
- 4) Estimate population sex- and age- structure.

Funding for this study was provided by the U.S. Fish and Wildlife Service. I gratefully acknowledge the contributions of the Montana Cooperative Wildlife Research Unit, Montana Department of Fish, Wildlife and Parks, and Glacier National Park.

## STUDY AREA

This study was conducted in the valley of the North Fork of the Flathead river in northwestern Montana and southeastern British Columbia. The study area extended from Anaconda creek in Glacier National Park northward to 30 km beyond the Canadian border.

The North Fork valley bottom varies from 4-10 km in width and rises from 1024 m elevation in the south to 1375 m in the northern part of the study area. Peaks of the Whitefish range form the western border of the valley, and the Livingston range defines the eastern border.

Land east of the North Fork of the Flathead river lies in Glacier National Park. West of the river, land is a mosaic of



Flathead National Forest and private property. North of the international border, land on both sides of the river is primarily under British Columbia provincial ownership.

Mean temperature ranges from -9 C in January to 16 C in July (Singer 1979). Snow normally covers the study area from mid-November through mid-April.

Dense lodgepole pine (*Pinus contorta*) forests dominate most of the North Fork valley, but sub-alpine fir (*Abies lasiocarpa*), spruce (*Picea* spp.), western larch (*Larix occidentalis*), and Douglas-fir (*Pseudotsuga menziesii*) communities exist throughout the valley. Abundant meadows and riparian areas are dispersed within the study area. Detailed descriptions of vegetation communities in this area have been provided by Habeck (1970), Jenkins (1985), and Kraemer (1989).

## METHODS

### Trapping

I selected 3 white-tailed deer wintering areas in Glacier National Park for trapping: Kintla Lake, Bowman Road, and the Sullivan Meadow area. These 3 winter ranges provided a northern, central, and southern sample of deer within the area inhabited by wolves. Deer were trapped from the Kintla Lake, Sullivan Meadow, and Bowman Road winter ranges from 21 January through 31 March 1990, and from 26 November 1990 to 26 February 1991.

Deer were trapped with modified elk-sized Clover traps (described by Thompson et al. 1989) or standard Clover traps (Clover 1956). All traps were baited with alfalfa hay (certified to be free of noxious weeds). Female white-tailed deer were manually restrained and instrumented with a radio transmitter (MOD-500, Telonics, Inc., Mesa, Ariz.) with a mortality sensor (4-hr delay). When does >1 yr old were captured, I extracted a canine tooth (I<sub>4</sub>) for use in age-determination. Deer were released following tooth extraction. Male white-tailed deer and all mule deer were released without being handled.

### Mortality

I tried to monitor activity signals of all radio-collared deer at least once daily. When a radio signal indicated that a deer had not moved in >4 hrs, I carefully approached the animal on the ground and performed a post-mortem examination to determine cause of death (O'Gara 1978, Wobeser and Spraker 1980). When predation had occurred, I recorded kill and chase information, and when possible, attempted to establish the pre-mortality condition of the animal via analysis of femur marrow, kidney fat index, and description of other vital organs (Thorne et al. 1982). If near-total consumption of the carcass made it impossible to ascertain if the deer was killed or was scavenged soon after death, I attributed cause of death to what I considered the most likely scenario and labeled the death a "probable" predation. Seasonal and yearly cause-specific mortality rates will be computed with program MICROMORT (version



1.3, Heisey 1987) and methods of Heisey and Fuller (1985) upon completion of field work.

### Seasonal Distribution

To identify key areas of seasonal use and document movement patterns, I attempted to locate all deer weekly by triangulating at least 3 strong radio bearings. I plotted radio bearings on USGS (1:24,000) or Energy, Mines and Resources Canada (1:50,000) topographic maps, and selected a location either at the center of the smallest triangle defined by 3 or more signal azimuths, or at the intersection of 2 such triangles. I divided locations into 6 categories of precision (<1 ha, 1-3 ha, 3-6 ha, 6-12 ha, 12-25 ha, or >25 ha) based on size of the triangle, or "error polygon." Variable topography and lack of an extensive road network within the study area frequently inhibited my ability to get close-range, line-of-sight signal fixes. Consequently, precise triangulations were often difficult to obtain. Only locations with error polygons of <25 ha were used in calculation of seasonal ranges. If I could not locate a deer from the ground, I located it from a Cessna 180 airplane when possible.

I computed minimum convex polygon and 95% harmonic mean (25 grid cell) range for winter and summer ranges of each radio-collared deer (McPAAL ver. 1.2, Stuwe and Blohowiak nd.). To estimate migration distances, I calculated the straight-line distance between the approximate center of each deer's winter and summer range.

### Index of Population Abundance

I initiated an intensive pellet-group sampling scheme to monitor trend of the white-tailed deer population over time. Methods were based on non-permanent 1.8 m-radius plots and variability estimates of Tucker (Thesis in prep). Pellet group sampling began in April concurrent with disappearance of snow cover. In 1990 I counted pellet groups in 80 plots on 7.5 pairs of transects (n = 600 plots) in areas with lower density of deer (Fig. 3), and 40 plots in 20 0.25 km<sup>2</sup> blocks (n = 800 plots) in areas with high density of deer (Kintla Lake/Starvation Ridge area, Bowman Road/Akokala area, and Sullivan Meadow). Based on variance estimates obtained from 1990 data, I will revise this sampling scheme to enable future researchers to detect a 20% change in white-tailed deer population numbers (Confidence Limit = 0.90).

### Population Sex- and Age- structure

During spring green-up, large numbers of white-tailed deer gather in fields along the North Fork Road to take advantage of the new grass shoots. One hour before sunset, from mid-April to mid-May, 1990 and 1991, I drove from 1.6 km south of Coal Creek (mile marker 24) to Polebridge (mile marker 32) and searched for deer. All deer were counted and observed with 10x binoculars. If possible, deer were classified as adult (>1 yr-old) males or females, or fawns (<1 yr-old). Upon completion of the field



work, age-structure estimated from roadside counts will be compared to age-distribution of does killed by hunters in November and December of the previous year.

## RESULTS AND DISCUSSION

From January through March 1990, 50 white-tailed deer and 11 mule deer were captured. Twenty-three white-tailed deer females were fitted with radio collars. During winter 1990-1991, 54 white-tailed deer were captured and 15 females were radio-collared. In March 1990 1 radio transmitter apparently ceased functioning.

### Mortalities

Between February 1990 and March 1991, 7 radio-collared deer were killed. Wolves and humans each killed 2 radio-collared deer, and a bear, mountain lion, and coyote each killed 1 (Table 1). Wolves killed a 2.5 yr-old female on 13 March 1990, and probably killed a 10 yr-old female on 4 June 1990. Because the carcass of the second deer was heavily scavenged by black bears, I was unable to determine if wolves had actually killed the deer or only scavenged the carcass. Bears probably killed an old (11-13 yrs) radio-collared deer on 13 June 1991. Lack of remains prevented positive determination of whether the deer was killed by bears or scavenged soon after death. Humans were responsible for the death of a deer on 26 June 1990, and another during hunting season on 5 November 1990. A mountain lion killed a deer on 31 January 1991, and coyotes killed a fawn (approx. 8 months old) on 13 February 1991.

Predators killed 3 radio-collared deer during winter and 2 during the fawning period in June, but killed none during summer or autumn (Table 1). Both deer killed during the June fawning period were very old (Table 1). Because older deer may be in weakened physical state after a long winter and the increased physical demands of pregnancy, they are likely to very vulnerable to predation during the fawning period. Lack of predation on adults during summer and autumn months suggests that predators may be relying on an abundance of fawns during this period (Van Ballenberghe et al. 1975, Fritts and Mech 1981, Nelson and Mech 1986).

### Seasonal Distribution

Radio-collared white-tailed deer wintered in 3 areas within the range inhabited by wolves in western Glacier National Park: the Kintla Lake area, the Polebridge/Bowman Lake area, and the Sullivan Meadow area. Females migrated from winter ranges during May and travelled an average of 12.0 km ( $n = 21$ ,  $SD = 11.6$ ) to summer ranges (Table 2). One deer migrated 40 km from winter range to summer range, but 4 deer did not migrate from wintering areas and lived in the same area year-round (Table 2). Deer began migrating from summer ranges as early as 26 August and arrived on winter ranges as early as 16 October. All deer



migrated from their summer ranges by mid-December. Of the 13 deer that migrated from summer ranges to winter ranges in 1990, 6 used intermediate "transitional" ranges for > 2 months, probably because of the lack of snow prior to late November.

Seasonal distribution of white-tailed deer in this area does not appear to be influenced by the presence of wolves. During summer, 5 radio-collared deer spent the summer within 1 km of an active wolf den in Sullivan Meadow. Wolves were in the meadow area continually from mid-April through December 1990, but the deer seemed unaffected by the presence of wolves and remained in the area.

#### Population Index

In April and May 1990, 1,667 deer pellet groups were counted in 1,400 1.8m-radius non-permanent plots ( $\bar{x}$  = 1.2 groups/plot, SD = 2.0). This pellet sampling index will be revised and continued in spring 1991.

#### Sex- and Age- Composition

I counted 1173 white-tailed deer in 11 evenings during April and May 1990 ( $\bar{x}$  = 106.6 deer/evening, SD = 29.2). I categorized 102 deer as adult males (15%), 429 as adult females (62%), and 155 as fawns (23%); 487 deer could not be classified. Roadside surveys will continue in Spring 1991.

#### CONCLUSION

Results presented in this paper are from the first year of a 2-year study. I feel strongly that it is premature to attempt to make specific conclusions based on these preliminary data. Consequently, I have refrained from making specific conclusions and have provided only a minimal and general discussion of results. Field work for this study will continue through the end of August 1991. After this time I will offer a more detailed discussion and attempt to make reliable conclusions.

#### LITERATURE CITED

- Ballard, W. B., J. S. Whitman, and C. L. Gardner. 1987. Ecology of an exploited wolf population in south-central Alaska. *Wildl. Monogr.* 98. 54pp.
- Bath, A. J. 1987. Attitudes of various interest groups in Wyoming toward wolf reintroduction in Yellowstone National Park. M. S. Thesis. Univ. Wyoming, Laramie. 122 pp.
- \_\_\_\_\_, and T. Buchanan. 1989. Attitudes of interest groups in Wyoming toward wolf restoration in Yellowstone National Park. *Wildl. Soc. Bull.* 17:519-525.
- Boyd, D. K., R. R. Ream, M. W. Fairchild, and D. H. Pletscher. in manuscript. Prey characteristics of colonizing wolves in



the Glacier National Park area.

- Clover, M. R. 1956. Single-gate deer trap. Calif. Fish and Game 42:199-201.
- Day, G. L. 1981. The status and distribution of wolves in the northern Rocky Mountains of the United States. M. S. Thesis. Univ. Montana, Missoula. 130 pp.
- Fritts, S. H., and L. D. Mech. 1981. Dynamics, movements, and feeding ecology of a newly protected wolf population in northwestern Minnesota. Wildl. Monogr. 80. 79 pp.
- Habeck, J. R. 1970. The vegetation of Glacier National Park. Natl. Park Serv. and Univ. Montana. 123 pp.
- Heisey, D. M. 1987. MICROMORT user's guide. NH Analytical Software, St. Paul, Minn. 10 pp.
- \_\_\_\_\_, and T. K. Fuller. 1985. Evaluation of survival and cause-specific mortality rates using telemetry data. J. Wildl. Manage. 49:668-674.
- Jenkins, K. J. 1985. Winter habitat and niche relationships of sympatric cervids along the North Fork of the Flathead River, Montana. Ph. D. Dissertation. Univ. Idaho, Moscow. 183 pp.
- Kellert, S. R. 1985. The public and the timber wolf in Minnesota. Yale Univ., New Haven, CT. 175 pp.
- Krahmer, R. W. 1989. Seasonal habitat relationships of white-tailed deer in northwestern Montana. M. S. Thesis, Univ. of Montana, Missoula. 104 pp.
- McNaught, D. A. 1987. Wolves in Yellowstone National Park? - - Park visitors respond. Wildl. Soc. Bull. 15:518-521.
- Mech, L. D. 1966. The wolves of Isle Royale. U. S. Natl. Park Serv. Fauna Ser. 7. 210 pp.
- Messier, F., and M. Crete. 1985. Moose-wolf dynamics and the natural regulation of moose populations. Oecologia 65:503-512.
- Murie, A. 1944. The wolves of Mount McKinley. U. S. Natl. Park Serv. Fauna Ser. 5. 238 pp.
- Nelson, M. E., and L. D. Mech. 1986. Mortality of white-tailed deer in northeastern Minnesota. J. Wildl. Manage. 50:691-698.



- O'Gara, B. W. 1978. Differential characteristics of predator kills. Pages 380-393 in Proc. 8th Biennial Pronghorn Antelope Workshop, Jasper, Alberta.
- Ream, R. R., M. W. Fairchild, D. K. Boyd, and D. H. Pletscher. 1987. Wolf monitoring and research in and adjacent to Glacier National Park. Section 6 Final Report. 41 pp.
- , -----, -----, and A. J. Blakesly. 1989. First wolf den in western U. S. recent history. Northwestern Naturalist 70:39-40.
- Singer, F. J. 1979. Habitat partitioning and wildfire relationships of cervids in Glacier National Park, Montana. J. Wildl. Manage. 42:437-444.
- Telonics Telemetry-Electronics Consultants, Mesa, Ariz.
- Thompson, M. J., R.E. Henderson, T. O. Lemke, and B. A. Sterling. 1989. Evaluation of a collapsible clover trap for elk. Wildl. Soc. Bull. 17:287-290.
- Thorne, E. T., N. K. Kingston, W. R. Jolley, and R. C. Bergstrom. 1982. Diseases of wildlife in Wyoming, 2nd Ed. Wyoming Game and Fish Dept. 353 pp.
- Tucker, P., and D. H. Pletscher. 1989. Attitudes of hunters and residents toward wolves in northwestern Montana. Wildl. Soc. Bull. 17:509-514.
- Van Ballenberghe, V., A. W. Erickson, and D. Byman. 1975. Ecology of the timber wolf in northeastern Minnesota. Wildl. Monogr. 43. 43 pp.
- Wobeser, G. A., and T. R. Spraker. 1980. Post-mortem examination. Pages 89-98 in S. D. Schemnitz, ed. Wildlife management techniques manual. 4th ed. The Wildl. Soc., Washington, D. C.



Table 1. Cause-specific mortality of radio-collared female white-tailed deer in the North Fork drainage of the Flathead River, February 1990 - March 1991.

Cause	Predator	Date	Age of Deer (yrs)
Predation	Wolves	03/13/90	2.5
Probable Predation	Wolves	06/04/90	10.0
Probable Predation	Bear	06/13/90	11-13
Accident <sup>1</sup>	Humans	06/26/90	8.0
Hunting	Human	11/05/90	6.5
Predation	Mountain Lion	01/31/91	5.5
Predation	Coyote	02/13/91	0.75

1 Killed by accident during translocation attempt in Glacier National Park.



Table 2. Seasonal distribution and migration distances of white-tailed deer radio-collared in Glacier National Park (GNP) in winter 1990.

ID#	Winter Range in GNP	Summer Range	Migration Dist. (km)
101	Kintla Lake	Kintla Lake	0
102	Kintla Lake	Colts Cr., Pvt <sub>1</sub>	11
103	Kintla Lake	Couldrey, Cr., BC	21
105	Kintla Lake	Kintla Lake	0
106	Kintla Lake	Ford Cr., GNP	5
107	Kintla Lake	Harvey Cr., BC	40
108	Kintla Lake	8 km S. Polebridge, Pvt	27
109	Kintla Lake	Couldrey Cr., BC	21
110	Polebridge	Polebridge, Pvt	1
111	Sullivan Meadow	Dutch Creek, GNP	8
112	Sullivan Meadow	Big Prairie, GNP	16
113	Sullivan Meadow	Mid. Quartz Lake, GNP	14
114	Sullivan Meadow	Sullivan Meadow	0
115	Sullivan Meadow	Sullivan Meadow	0
116	Sullivan Meadow	Sullivan Meadow	2
117	Sullivan Meadow	Mud Lake, GNP	7
118	Sullivan Meadow	Hay Creek, FNF <sub>2</sub>	23
119	Sullivan Meadow	Logging Cr., GNP	3
120	Sullivan Meadow	Hidden Meadow, GNP	7
121	Sullivan Meadow	Tepee Lake, FNF	31
123	Sullivan Meadow	Anaconda Creek, GNP	16

1 Pvt = Private property

2 FNF = Flathead National Forest



## The Significant of Disease in Wildlife Populations

This talk is oriented toward wildlife biologists organizing and conducting population ecology studies. It is for those of you who have asked the question, "Do I need to add disease investigations to my population study in order to be complete?". The answer is not always "yes". Hopefully this presentation will increase your familiarity with diseases in wildlife populations and help you begin to answer this question.

The word "disease" is very general. It is defined as any condition that impairs the normal structure or function of an individual or population. Disease can include infectious agents, parasites, nutritional problems, environmental toxins, cancers and physical damage - anything that causes "dis-ease".

Disease ecology is one of the many bridges connecting wildlife ecology and veterinary science. Disease ecology looks at how diseases influence, and are influenced by, the dynamics of animal populations. It relates individual physiology and disease to population health and condition. I like to consider disease ecology as being more of a population issue in which the research biologist and manager can contribute greatly utilizing veterinary skills and perspectives for data collection.

Infectious diseases are very dynamic in wildlife populations. We often perceive them as being in some quiescent, stagnant form, but they are ever changing. One example is prevalence (number of animals infected) of a disease, which can fluctuate over the years<sup>1</sup>. Infectious diseases also exert continuous pressure on populations. We generally view them negatively, but they often play a very positive role in population regulation and adaption<sup>2</sup>.

Their impact is very complex, being influenced by factors such as weather, predators and intra- and interspecific competition. Weather and other environmental factors can affect the abundance of disease organisms. Predators selecting for compromised prey can serve as regulators of disease populations. Competition and nutrition can contribute stress factors increasing the population's susceptibility to disease.

There is constant interaction between hosts and infectious agents. Hosts are continuously applying pressure on the disease agent in an effort to fight off infection. This pressure, in turn, influences the infecting organism to change in order to be more effective or less evident to the host. Such a "tug-of-war" produces diseases that are continuously evolving (sometimes into new diseases) and populations with changing disease resistance<sup>4</sup>.

Some diseases, such as the footworm (*Wehrdikmansin cervipedis*) and abdominal worm (*Setaria yehi*) of elk, have evolved with the host into an almost symbiotic relationship resulting in little damage to either parasite or host. Such diseases, obviously, may not be cost-effective for a field biologist to include into ecological studies.

From a practical perspective, the above relationships between disease agents, host and environment suggests that there are no easy answers and that data may not always provide conclusive results. It also supports the need for disease investigations so we can clarify these complex relationships.

There are a variety of roles that diseases play in wildlife populations. The following are some brief examples of these roles.

### Diseases can directly limit populations by being life threatening.

Canine parvovirus and feline panleukopenia can easily cause death to individuals



previously unexposed (and therefore unprotected). Losses from these diseases commonly occur in young animals near the time of weaning when it is difficult to identify the occurrence of mortalities in free-ranging wildlife and even more difficult identifying the cause of mortalities.

Rabies has decimated entire wolf packs in coastal Alaska<sup>5</sup>, but does not appear to play a similar role in interior Alaska or in Minnesota wolf populations. The reason for this difference is unknown.

Bluetongue occurs in livestock and wildlife over much of the United States. In the Rocky Mountain region, this disease is often neglected in wildlife studies. Yet in 1976, bluetongue killed a minimum of 1400 antelope and deer in eastern Wyoming<sup>6</sup> emphasizing the importance of this disease in wild ruminants.

Diseases can directly limit populations by causing reproductive failure. *Brucella abortus* in bison and elk of the Yellowstone ecosystem is a notable example of this role. The major symptoms of brucellosis in cows are abortions, premature births and birth of non-viable calves<sup>6</sup>. Without disease investigations, these losses due to reproductive failure as well as those causing mortality of young could easily be overlooked or mistaken for predation, starvation or other factors.

Diseases can indirectly limit populations.

As stated above, disease agents continually exert pressure. Their presence is one more additional stress that can be as significant as many environmental factors we frequently consider in our wildlife populations. Dr. Terry Spraker describes the scenario of a dog swimming in a pond. Place a small weight on his back and he will continue to swim strong and high in the water. But with each additional weight, he will be slower and lower in the water. Finally, there will be a single weight that keeps the dog from reaching the other shore. It is not the final weight which was most significant for the dog's loss, but an accumulation of weights, just as our wildlife populations have an accumulation of stresses placed upon them including diseases.

Winter ticks are a good example of an additional stress to wildlife. They can be a serious parasite for moose and elk causing anemia due to blood loss and hypothermia due to hair loss.

Diseases can support one species in competing against other species.

Dramatic examples of this role are the arterial worm (*Elaeophora schneideri*) and the meningeal worm (*Parelaphostrongylus tenuis*). The mule deer benefits from having *E. schneideri* as a parasite. This parasite reaches maturity within carotid arteries of the mule deer, produces microfilariae in the bloodstream which are picked up by horseflies and transmitted to other ungulates. This parasite is fatal to many abnormal hosts such as elk and moose resulting in blockage of arteries supplying the head, blindness, loss of body parts and death. In areas of Arizona and New Mexico where mule deer, horseflies and the arterial worm are abundant, elk populations are limited<sup>7</sup>.

The meningeal worm benefits its normal host, white-tailed deer, and is devastating for many other ruminants including elk, moose and mule deer. It exists primarily in the eastern United States. Adult parasites reside in the membranes (meninges) covering the brain, produce eggs that hatch into larvae. These larvae enter the bloodstream, travel to the lungs,

are c  
snails  
these  
spina

State  
disea

Wild

withi  
such  
meth  
stres:  
modi

of th  
share  
varic  
in lo  
wildl

study  
Leop  
prob

scier  
the i  
Fort

are 1  
samj  
train

with  
curr  
prev  
som

the  
inex  
prog

han  
reas



are coughed up and swallowed, and exit with the feces. The larvae then develop within land snails into infective stages which are picked up by grazing individuals. In abnormal hosts, these larvae migrate to the brain also, but wander throughout the tissue of the brain and spinal cord causing life-threatening damage<sup>7</sup>.

The meningeal worm prevents successful elk reintroductions into the eastern United States. It is also a strong example of why we must be careful transporting wildlife (and their diseases), in this case, from eastern to western parts of the country.

#### Wildlife diseases can be indicators for the biologist.

Lungworm/pneumonia complex in bighorn sheep is usually an indicator of stress within the population. With outbreaks occurring, other stress factors are often identified, such as human activity, hunting, and high population density. Researchers have developed methods to monitor a stress hormone, cortisol, within sheep populations and can identify stressed populations before disease outbreaks occur. Once identified, wildlife managers can modify factors to reduce stress and prevent all age die-offs (T. Spraker, pers. comm.).

Some diseases are more prevalent in domestic animals than in wildlife. The presence of these diseases in wildlife populations can suggest that the population is utilizing habitat shared with domestic animals - information often valuable to the biologist. Prevalence of various diseases within the wildlife population can be compared to known prevalence rates in local livestock. Once identified, modes of transmission and disease influences for both wildlife and domestic animals should be considered.

These are just a few of the roles of disease in wildlife populations.

As we decide whether or not to incorporate disease investigations into a population study, it is worth asking, "Why has there been so little done in disease ecology?". Aldo Leopold stated almost 60 years ago, "The role of disease in wildlife conservation has probably been radically underestimated." This is still true today.

Traditionally, there has been a separation between wildlife biology and veterinary science. Wildlife biology has worked with populations collecting data on factors outside of the individual. Veterinary science has focused on the individual and its internal activity. Fortunately, these two fields are interacting more all the time.

Blood collecting, necropsies, and other data collection skills for disease investigations are not routinely taught as part of university wildlife programs. Transport and handling of samples and interpretation of results can be equally obscure without proper consultation and training.

Once data is collected, there is not a great deal that the manager feels he can do with it. Eradication of disease in free-ranging populations appears impossible with our current technology. We don't even have a clear understanding of how to reduce the prevalence of most diseases. But disease studies do contribute to eventual eradication of some diseases and increase our understanding of population dynamics.

With limited finances and manpower the researcher and manager must also consider the cost/benefit of every investigation. Disease studies can be very costly, or at times, very inexpensive. Logistically, sample collections are easily incorporated into animal handling programs for most field conditions.

Moral obligations and ethical values should also be considered. As we capture and handle wildlife, we have a responsibility to gather as much information as possible, within reason. If diseases appear to play a significant role, scientific investigation of population



dynamics would be incomplete without including disease aspects. Also with the many wildlife diseases affecting human health and livestock, it is the field biologist who can play a significant role in understanding and eradicating these diseases.

Lastly, and most significantly, diseases are often neglected in wildlife studies because there is so little known about the disease ecology of most diseases. Often the questions are not even developed, let alone acquiring answers. Again it is the field biologist working with veterinary perspectives that can provide insight and answers.

There are a variety of resources for the researcher and manager for assistance in developing and conducting disease investigations. These include:

- Biologists with strong backgrounds in wildlife diseases
- State diagnostic laboratories and wildlife research labs
- Members of the Wildlife Disease Association that are publishing papers on the species you are studying
- Wildlife veterinarians that combine veterinary science with population perspectives and a practical understanding of field conditions.

### References Cited

1. Webb, P.A. et.al.1987. Epizootic vesicular stomatitis in Colorado, 1982: Some observations on the possible role of wildlife populations in an epizootic maintenance cycle. *J. Wildl. Dis.* 23(2):192-198.
2. Friend, M. 1976. Wildlife diseases: Philosophical considerations. In: Proceedings of the Third International Wildlife Disease Conference. L.A. Page,ed. New York Plenum Press. pp. 7-43.
3. Holmes, J.C.1982. Impact of infectious disease agents on the population growth and geographical distribution of animals. In: Population Biology of Infectious Diseases. R. M. Anderson and R.M. May, eds. New York Springer-Verlag. pp. 37-51.
4. O'Brien, S.J. and J.F. Evermann. 1988. Interactive influence of infectious disease and genetic diversity in natural populations. *Trends in Ecology and Evolution* 3(10):254-259.
5. Zarnke, R.L. and W. B. Ballard.1987. Serologic survey for selected microbial pathogens of wolves in Alaska, 1975-1982. *J. Wildl. Dis.* 23(1):77-85.
6. Thorne, E.T. et.al.,ed.1982. Diseases of Wildlife in Wyoming. Wyoming Game and Fish Department.
7. Kistner, T.P.1982. Diseases and parasites. In: Elk of North America, Ecology and Management. J.W. Thomas and D.E. Toweill, eds. pp.181-217.

Mark R. Johnson, DVM  
P.O. Box 883  
Gardiner, MT 59030

BLAC

WAYN  
Libb

TIMO  
Libb

Abst:  
amer  
Data

(HD  
appro  
calcu

appro  
had  
table

repor  
indio

great  
was  
surv

colla  
Annua  
from

calcu  
for  
lower

to m  
acce

INTR

popu  
of F

days  
55,0  
hunt  
bear  
decl

the  
hunt

limi  
to en

Wadd  
great  
short  
weeks  
blac



## BLACK BEAR SURVIVAL RATES IN NORTHWEST MONTANA

WAYNE F. KASWORM, U.S. Fish and Wildlife Service, 475 Fish Hatchery Road, Libby, MT 59923

TIMOTHY J. THIER, U.S. Fish and Wildlife Service, 475 Fish Hatchery Road, Libby, MT 59923

**Abstract:** Survival estimates were calculated for black bears (*Ursus americanus*)  $\geq$  4 years old for the period 1983-1990 in northwest Montana. Data was examined from several studies on black bears in 3 Hunting Districts (HD 100, HD 104, and HD 121) using 3 different approaches. In the first approach a computer model (MICROMORT; Heisey and Fuller 1985) was used to calculate survival for radio collared bears only ( $N = 45$ ). In the second approach, life tables were calculated for all black bears  $\geq$  4 years old that had been captured, tagged, and released ( $N = 189$ ). In the third approach, life tables were calculated for all bears  $\geq$  4 years old that had been killed and reported by hunters ( $N = 617$ ). Capture information and radio monitoring indicated that the age of first successful reproduction was 6 years old or greater, mean litter size was 1.7 cubs, and the mean interval between litters was 3.5 years. Based on this information, a minimum sustainable annual survival rate of 0.89 was estimated. Composite annual survival rates for radio collared bears ranged from 0.49-0.77 for males and 0.68-0.89 for females. Annual survival rates calculated from life tables for captured bears ranged from 0.70-0.81 for males and 0.72-0.84 for females. Annual survival rates calculated from life tables for bears killed by hunters ranged from 0.69-0.75 for males and 0.73-0.74 for females. All analyses indicated survival rates lower than the minimum sustainable rate of 0.89. Management strategies related to mortality (e.g. hunting season timing, hunting season length, hunter access, and enforcement) were discussed.

### INTRODUCTION

Black bear hunting in northwest Montana has shown a dramatic increase in popularity since the early 1970's. In 1971, Region 1 of the Montana Department of Fish, Wildlife, and Parks (MDFWP) recorded about 19,000 black bear hunter days and approximately 650 bears killed. By 1979, hunter days had increased to 55,000 with almost 1,000 bears harvested. With this increase in harvest and hunting pressure, MDFWP biologists became concerned about trends in overall bear numbers. Interviews with hunters and residents indicated a general decline in bear sightings and nuisance bear complaints.

An average of 62% of the annual harvest from 1978-1980 occurred during the spring hunting season, with most during May. Black bears are vulnerable to hunting in spring because snow cover forces animals searching for food into limited areas of early green-up. Females (particularly females with cubs) tend to emerge from dens later in the spring than males (Hugie 1982, Beecham 1980, Waddel and Brown 1984). In an effort to reduce the overall harvest and afford greater protection to reproductively active females, MDFWP biologists shortened the spring season in districts 100, 101, 103, and 104 to only 2 weeks in late April beginning in 1981. After 3 years of curtailed harvest, black bear populations appeared to increase. In response, the spring season



was lengthened to 4 weeks in 1984 (15 May ending date). This paper seeks to compare and evaluate mortality rates under this hunting regime using data from hunting districts 100, 104, and 121.

## STUDY AREA

Habitat use and population characteristics of black and grizzly bears were studied from 1983 to 1990 in the Cabinet Mountains and the Yaak River drainage of northwest Montana (48° N, 116° W). The study area encompassed approximately 7,800 km<sup>2</sup> and is bisected by the Kootenai River, with the Cabinet Mountains to the south and the Yaak River area to the north (Figure 1). Approximately 90% of the study area was on public land administered by the Kootenai and Panhandle National Forests. The Cabinet Mountains Wilderness Area encompassed 381 km<sup>2</sup> of the study area at higher elevations of the Cabinet Mountains.

Elevations on the study area ranged from 664 m along the Kootenai River to 2,664 m at Snowshoe Peak. Weather is dominated by a Pacific maritime climate characterized by short, warm summers and heavy, wet winter snowfalls. South and west slopes at lower elevations supported stands of ponderosa pine (*Pinus ponderosa*) and Douglas-fir (*Pseudotsuga menziesii*). Grand fir (*Abies grandis*), western red cedar (*Thuja plicata*), and western hemlock (*Tsuga heterophylla*) dominated the lower elevation moist sites. Mixed stands of subalpine fir (*Abies lasiocarpa*), spruce (*Picea engelmannii*), and mountain hemlock (*Tsuga mertensiana*) were predominant above 1,500 m. Lodgepole pine (*Pinus contorta*) dominated large areas at mid- and upper elevations, especially north of the Kootenai River. Mixed stands of coniferous and deciduous trees were interspersed with riparian shrubfields and wet meadows along the major rivers. Huckleberry (*Vaccinium* spp.), an important food for black and grizzly bears, was a common component in the understory. The occurrence of huckleberry and other berry-producing shrubs were largely a result of wildfires that occurred between 1910 and 1929, and also from timber harvesting. Effective fire suppression since then has virtually eliminated wildfire as a natural force in creating and maintaining berry-producing shrubfields.

The southern portion of the study area was characterized by high, precipitous peaks with steep slopes. The northern portion was characterized by mountains that were lower in elevation, had gentler slopes, and were forest-covered. Contemporary resource use includes mineral exploration and extraction, timber harvesting, and recreation.

## METHODS

### Capture and Marking

Bears were captured with leg-hold snares following the techniques described by Johnson and Pelton (1980). The snares are manufactured by Aldrich Snare Co. (Clallam Bay, WA) and consist of 0.25 inch wound steel aircraft cable. All bears were immobilized with a drug mixture of Ketaset (ketamine hydrochloride) and Rompun (xylazine hydrochloride). Drugs were administered intramuscularly with either a syringe mounted on a pole or a Palmer Cap-Chur gun. Immobilized bears were measured, weighed, tattooed, and a first premolar tooth extracted for age determination (Stoneberg and Jonkel 1966).






Fig. 1. Map of study area in northwest Montana.



Each bear was marked with an individually numbered ear tag in each ear. Attached to each ear tag was a 4 X 13 cm streamer of armatite. The color of the streamers varied by species and the year in which the animal was captured. Only adult black bears ( $\geq 4.0$  years old) were fitted with radio collars. Motion-sensitive collars manufactured by Telonics (Mesa, Arizona) were used on most bears. To prevent permanent attachment, a canvas spacer was placed in the collars that was designed to separate in 2-3 years, (Hellgren et al. 1988).

Trapping efforts were conducted during the spring (prior to mid-July) from 1983-1987 and 1989-1990. Fall trapping efforts (early September to mid-October) were conducted in 1984, 1986, 1987, 1989, and 1990. Trap sites were usually located within 200 m of a road to allow vehicle access. Many trap sites were behind closed roads. One or two 2-person crews checked the snares daily. Bait consisted primarily of road-killed deer, with lesser amounts of beaver carcasses, elk, moose, and meat scraps from processing plants.

#### Radio Monitoring

Weekly aerial radio locations were obtained (weather permitting) on instrumented black bears each week during the 7-8 month period in which they were active. In addition, efforts were made to obtain as many ground locations as possible, usually by triangulating from a vehicle. Bear activity was assessed each time a location was obtained. Collars that were inactive for unusual periods of time were approached from the ground and a determination was made of the bear's fate. All specific locations were plotted on USGS topographic maps (1:24,000) and recorded by Universal Transverse Mercator Coordinates (UTM's). Distance from radio locations to the nearest open road and trail were measured on maps to 30-m accuracy. Closed roads (closed to motorized vehicles) were considered trails in the analysis.

#### Survival Estimation

The MICROMORT computer software package was used to calculate survival and cause-specific mortality rates from the radio telemetry information (Heisey and Fuller 1985). This method required the assumptions that marked individuals were representative of the population and that individuals within sex and age classes had independent and equal probabilities of survival. Only radio-collared bears were used in determining survival rates.

Mortality dates were established from hunter harvest information and estimates of dates based on radio telemetry, collar retrieval, and mortality site inspection. Radio failure or suspected mortality dates were estimated by using the date of the last radio location when the animal was known to be alive. Survival rates included suspected mortalities from radio-collared bears that disappeared with less than 2.5 years of elapsed time on their transmitters.

Seasons were defined on the basis of spring and fall hunting season dates. These periods were (1) spring, 15 April to 31 May, (47 days) (2) summer, 1 June to 31 August, (92 days); (3) autumn, 1 September to 30 November, (91 days); (4) winter, 1 December to 14 April (135 days).

Life table analysis (Downing 1980) was applied to samples of captured and hunter harvested black bears. Means were calculated from sex and age specific survival rates of bears  $\geq 4$  years-old. Hunter harvest information from the mandatory black bear tooth return program was obtained from MDFWP records. It



was estimated that this sample represents about 70% of all harvested black bears and has been in effect since 1985 (Brown et al. 1987). Hunting district 103 was divided into 2 districts (103 and 104) in 1989. Harvest information referenced as district 104 includes all of district 103.

## RESULTS

### Reproduction

Black bear capture and radio monitoring data were examined to determine reproductive parameters for females in the study area. Ninety-eight captures of female black bears from 1983-1990 were analyzed to determine first age of reproduction. Incidence of estrous, lactation, cub presence, and nipple length were used in the analysis (Table 1). Lactation or cub presence was noted in only 1 of 27 bears aged 3-5 years-old. The 4 year-old bear that did show signs of lactation was radio collared and was never seen subsequently with cubs. Signs of estrous were noted in 15% ( $n = 27$ ) of the black bears aged 3-5 years-old. Five of eight 6 year-olds showed signs of estrous, were lactating, or had cubs present at capture. Fifty-six percent ( $n = 55$ ) of the captured black bears  $\geq 7$  years-old were in a similar reproductive state. Radio monitoring information indicated that no female bears first captured as 4 or 5 year-olds ( $n = 4$ ) produced cubs prior to the age of 6. Female black bears in our study area may come into estrous as early as 3 years-old, but few successfully produce cubs before the age of 6 years-old. Mean age of first reproduction may be  $> 6$  years-old.

Table 1. Reproductive state of female black bears captured in northwest Montana, 1983-1990.

Age Class	$n$	Estrous	Lactation or Cubs	Nipple Length $\geq 12$ mm ( $n^{\dagger}$ )
Cub-2.0	8	0	0	0 (5)
3.0	10	1	0	0 (7)
4.0	9	2	1	0 (6)
5.0	8	1	0	0 (6)
6.0	8	2	3	4 (5)
7.0	11	3	2	7 (9)
8.0	11	2	2	9 (10)
9.0	8	2	3	6 (7)
10.0-24.0	25	12	5	23 (23)

<sup>†</sup>Sample size for nipple length measurement.

We have postulated that nipple length is greater for bears that have produced cubs than for bears that have not produced cubs (Kasworm and Manley 1988). Nipple length measurements have been recorded on female bears captured since 1984 ( $n = 78$ ). None of the 24 captured black bears  $\leq 5$  years-old had nipple lengths  $\geq 12$  mm. Ninety-one percent of captured bears  $\geq 6$  years-old ( $n = 54$ ) had nipple lengths  $\geq 12$  mm. While this information does not



demonstrate a cause and effect relationship, it would seem unlikely that the dramatic correlation of increased nipple length and first reproduction could be completely coincidental. We believe this information supports our contention that little successful reproduction occurs on our study area before females reach 6 years of age.

Radio monitoring of 22 female black bears  $\geq 6$  years-old provided estimates of cub production and reproductive interval (Table 2). Fifty-nine female black bear-years of information was obtained from 1983-1990. Seventeen litters of cubs were known to have been produced during the study period. Two of those litters were known to have been lost because of the death of the mother before counts of cubs could be obtained. Fifteen known litters produced 25 cubs for a mean litter size of 1.7. Mean bear-years per litter was 3.5. Production of 25 cubs and the deduction of 2 bear-years from the total to account for the 2 known lost litters produced a calculated reproductive rate of 0.44. Assuming each of the 2 lost litters produced 2 cubs, the calculated reproductive rate would have been 0.49.

Table 2. Cub production of radio-collared female black bears  $\geq 6$  years-old in northwest Montana, 1984-1990.

	1984	1985	1986	1987	1988	1989	1990	Total
No. Cubs:	1	3	7	4	8	1	1	25
No. Litters:	1	1	7 <sup>1</sup>	2	4	1	1	17 <sup>1</sup>
No. Females Monitored:	5	12	18	12	8	2	2	59

<sup>1</sup>Includes 2 litters which were lost before counts of cubs were obtained.

#### Maximum Sustainable Mortality

A population model which estimates maximum sustainable mortality was developed by Bunnell and Tait (1980). This model assumed that the mortality rate is constant for all age-classes and that cubs die only if the mother dies. The mortality rate was balanced against the natality rate needed to produce a nondeclining population. Using an average litter size of 1.7, an interbirth interval of 3.5 years, and a first age of reproduction of 6 produces a maximum sustainable mortality rate of 11% from all causes. Survival rates less than 0.89 would cause this population to decline.

#### Survival of Radio Collared Black Bears

Forty-five black bears aged 4 to 21 years-old were radio collared in 3 study areas of northwest Montana during 1983-1990 (Appendix Table 1). Nineteen of these individuals were males that carried functional collars for 97-1,524 days. Twenty-six of the individuals were females that carried functional collars for 67-1,781 days. Eighteen instances of known mortality and 4 instances of suspected mortality were recorded over the 8 years of study. No mortality was detected in the sample during winter. Eight of 9 male deaths occurred during spring and 1 male died during summer. No male mortality was detected during autumn. Female mortality was distributed through 3 seasons



with 7 deaths occurring in spring and 3 each during both summer and autumn. Seasonal and annual survival estimates were calculated by sex for each hunting district within the study area (Table 3). No mortality was observed during winter. Interval survival rates for males were not different among districts (Z tests,  $P > 0.05$ ). Male survival rates were lowest during spring in all districts. Spring survival of females in district 104 was less than for females in district 121 ( $P < 0.05$ ). All other interval comparisons of females among districts were not different ( $P > 0.05$ ). Survival rates of females were lowest during spring in districts 100 and 104. Summer and autumn rates were equally low in district 121.

Table 3. Seasonal and annual survival rates of radio-collared black bears in northwest Montana, 1983-1990.

Sex	District	n	Radio Days	Season <sup>1</sup>			Annual (Deaths, 95% CI)
				Spring (Deaths, 95% CI)	Summer (Deaths, 95% CI)	Autumn (Deaths, 95% CI)	
<b>Male</b>							
	100	9	5,546	0.75 (4, 0.56-0.99)	1.00 (0)	1.00 (0)	0.75 (4, 0.56-0.99)
	104	8	4,708	0.77 (3, 0.57-1.00)	1.00 (0)	1.00 (0)	0.77 (3, 0.57-1.00)
	121	2	1,070	0.69 (1, 0.33-1.00)	0.71 (1, 0.36-1.00)	1.00 (0)	0.49 (2, 0.18-1.00)
	Total	19	11,324	0.75 (8, 0.61-0.91)	0.97 (1, 0.91-1.00)	1.00 (0)	0.73 (9, 0.59-0.90)
<b>Female</b>							
	100	9	4,634	0.82 (2, 0.62-1.00)	1.00 (0)	0.88 (2, 0.74-1.00)	0.72 (4, 0.52-1.00)
	104	12	8,038	0.75 (5, 0.58-0.96)	0.91 (2, 0.80-1.00)	1.00 (0)	0.68 (7, 0.52-0.91)
	121	5	5,969	1.00 (0)	0.94 (1, 0.83-1.00)	0.94 (1, 0.84-1.00)	0.89 (2, 0.75-1.00)
	Total	26	18,641	0.84 (7, 0.74-0.96)	0.94 (3, 0.88-1.00)	0.95 (3, 0.89-1.00)	0.75 (13, 0.64-0.88)

<sup>1</sup>No mortality was observed during winter.

Annual survival rates of males were not different ( $P = 0.516$ ) between districts 100 and 104 (0.75 and 0.77, respectively). Male annual survival rates in district 121 (0.49) were different ( $P < 0.001$ ) from districts 100 and 104. Similar relationships were observed among female black bears. Female annual survival rates were not different ( $P = 0.114$ ) between districts 100 and 104 (0.72 and 0.68, respectively). Annual survival rates of females in district 121 (0.89) were different ( $P < 0.001$ ) from districts 100 and 104. Pooled annual survival rates for males and females were 0.73 and 0.75, respectively.

#### Cause of Mortality

Nine male and 13 female mortalities occurred over the course of this study (Table 4). Twelve were reported hunter kills (8 male and 4 female), 2 were illegal kills (both females with cubs and outside of legal season), 1 was a natural mortality (female), 3 were of unknown origin (all female), and 4 were suspect mortalities based on the loss of the radio signal (1 male and 3



female). Forty-five percent of total mortality was unreported. Composite male and female annual mortality from all districts was 0.27 and 0.25 respectively. Hunting was the largest single source of mortality for males and females at 0.25 and 0.09 respectively. All other individual mortality sources were  $\leq$  0.06 each. Some of the mortality classified as unknown or suspect may have been related to hunting (e.g. wounding loss or failure to report kill).

Table 4. Pooled mortality rates by source for radio-collared black bears from hunting districts 100, 104, and 121 in northwest Montana, 1983-1990.

Mortality Source	Sex	Season <sup>1</sup>			
		Spring (Deaths, 95% CI)	Summer (Deaths, 95% CI)	Autumn (Deaths, 95% CI)	Annual (Deaths, 95% CI)
Hunting	Male	0.25 (8, 0.10-0.40)	0.00 (0)	0.00 (0)	0.25 (8, 0.10-0.40)
	Female	0.09 (4, 0.01-0.17)	0.00 (0)	0.00 (0)	0.09 (4, 0.01-0.17)
Illegal	Male	0.00 (0)	0.00 (0)	0.00 (0)	0.00 (0)
	Female	0.05 (2, 0.00-0.11)	0.00 (0)	0.00 (0)	0.05 (2, 0.00-0.11)
Natural	Male	0.00 (0)	0.00 (0)	0.00 (0)	0.00 (0)
	Female	0.00 (0)	0.02 (1, 0.00-0.06)	0.00 (0)	0.02 (1, 0.00-0.05)
Unknown	Male	0.00 (0)	0.00 (0)	0.00 (0)	0.00 (0)
	Female	0.00 (0)	0.00 (0)	0.05 (3, 0.00-0.11)	0.05 (3, 0.00-0.09)
Suspect	Male	0.00 (0)	0.03 (1, 0.00-0.09)	0.00 (0)	0.03 (1, 0.00-0.09)
	Female	0.02 (1, 0.00-0.07)	0.04 (2, 0.00-0.09)	0.00 (0)	0.06 (3, 0.00-0.12)
TOTAL	Male	0.25 (8, 0.09-0.39)	0.03 (1, 0.00-0.09)	0.00 (0)	0.27 (9, 0.10-0.41)
	Female	0.16 (7, 0.04-0.26)	0.06 (3, 0.00-0.12)	0.05 (3, 0.00-0.11)	0.25 (13, 0.12-0.36)

<sup>1</sup>No mortality was observed during winter.

#### Relations of Human Access to Mortality

Road and trail influences on black bears in hunting districts 104 and 121 were described by Kasworm and Manley (1990). This analysis involved distances from radio locations to the nearest open road or trail (trails include closed roads). This data set was further analyzed to determine the relations between mortality and black bear use of habitat near roads or trails. Only aerial locations were analyzed.

Mean distance to the nearest open road from 518 radio locations of bears killed during this study ( $x = 1,576$  m) was less (t-test,  $P < 0.001$ ) than the mean distance for 701 locations of bears still alive at the end of the study ( $x = 1,959$  m). Mean distance to the nearest trail from locations of bears killed during the study ( $x = 590$  m) was less (t-test,  $P = 0.043$ ) than the mean distance for bears still alive at the end of the study ( $x = 655$  m).



## Life Table Analysis of the Capture Sample

Trapping activities resulted in the capture of 189 black bears  $\geq 4.0$  years old in the 3 hunting districts during 1983-1990. Comparisons of male and female capture sample age distributions by hunting district showed no differences among years (Kruskal-Wallis,  $P > 0.177$ ). Yearly capture samples were pooled to form 6 samples of males and females from the 3 districts. Life table analysis was performed on each sample to estimate mean adult survival rates (Appendix Table 2). Mean survival rates for males  $\geq 4.0$  years old varied from 0.70-0.81 and mean survival rates for females varied from 0.72-0.84.

## Life Table Analysis of the Hunter Harvest

A mandatory tooth return program for successful Montana black bear hunters has been in effect since 1985. Hunter harvest from the 3 hunting districts was 617 black bears  $\geq 4.0$  years old from 1985-1989. Comparisons of male and female harvest sample age distributions by hunting district showed no differences among years (Kruskal-Wallis,  $P > 0.082$ ). Yearly harvest samples were pooled to form 6 samples of males and females from the 3 districts. Similar life table techniques as employed on the capture sample were applied to the harvest samples to estimate a mean adult survival rate for males and females  $\geq 4.0$  years-old (Appendix Table 3). Mean survival rates for males  $\geq 4.0$  years old varied from 0.69-0.75 and survival rates for females varied from 0.73-0.74.

## DISCUSSION

We calculated a natality rate of 0.49 and an age of first reproduction of 6 years-old for our study area. These values are similar to the natality rate of 0.57 and 6 years-old as a first age of reproduction for the Whitefish Range in northwest Montana (Jonkel and Cowan 1971). Studies in northern Idaho have indicated natality rates of 0.46 and a first age of reproduction at 5 years-old (Beecham 1980 in Stringham 1990). However, these values contrast sharply from central and eastern U.S. black bear populations which have reported natality rates of 1.04-1.45 and mean age of first reproduction of 3.6-6.3 years-old (Rogers 1987, Alt 1982 in Stringham 1990).

When reproductive rates of black bears and grizzly bears (*Ursus arctos horribilis*) are compared, the values are similar or lower for black bears. A study of grizzly bears on the Montana Rocky Mountain East Front (Aune and Kasworm 1989) indicated a natality rate of 0.85 and a mean age of first reproduction of 7 years-old. McLellan (1988) reported a natality rate of 0.86 and a mean age of first reproduction of 6 years-old for grizzly bears in the North Fork of the Flathead drainage in northwest Montana and southeast British Columbia. Considerable attention is currently given to reducing the mortality of grizzly bears, because of their low reproductive parameters. Similar concerns should be expressed for black bears if current population levels are to be maintained or increased.

The evidence presented indicates survival rates lower than the minimum sustainable, given the reproductive parameters of the population. At least one potential bias in our data involves the proximity of radio collared bears to roads. Black bears that used habitat closer to roads appear to have a higher



mortality rate than bears that used less densely roaded habitat. Since the majority of trapping occurred within 200 m of roads (both open and closed roads), it is possible that bears captured were more prone to being killed. A large proportion of the marked bears killed were harvested during the spring hunting season, even though the total kill was evenly divided between the spring and fall hunting seasons in recent years.

Another potential bias in our data involves the segment of the population sampled by radio telemetry. Only bears  $\geq 4$  years-old were collared and survival rates determined by telemetry represent only that age group. Lower survival of subadult than adult black bears has been reported (Bunnell and Tait 1985). However, given the magnitude of mortality in the 3 data sets, the low reproductive rates of the population, and the degree of access currently available, we believe that a more conservative approach is warranted in northwest Montana black bear management.

#### MANAGEMENT IMPLICATIONS

Management of the black bear in Montana has evolved from early status as a nuisance to that of a highly valued game animal. Hunting seasons in northwest Montana have also evolved from a year round season to the current 4 week spring season and the 12 week autumn season. Comparatively, deer and elk seasons are held only in the autumn for 5 weeks. It is illegal for hunters to take cubs or female bears with young and the use of bait or dogs in hunting black bears is prohibited.

Hunter harvest of black bears and associated unreported mortality appears to be the largest source of mortality. Several means of reducing hunter kill are available. Reducing the season length during either the spring or autumn or a combination of reductions could be applied.

Many resident and nonresident hunters in Montana receive a black bear tag as part of a combination license. Some of these hunters may not directly seek to kill a black bear, but harvest them incidental to other big game hunting. Reducing incidental take by selling tags only to hunters who are interested in black bear hunting may reduce harvest. Incidental take could be eliminated by closing the black bear season before the opening of general big game season.

Mortality quotas have been applied to black bear hunting in south central Montana. These quotas involve total mortality and female mortality. When one of these quotas is reached, the season closes on 48 hour notice. Hunting only by permit is another option to reduce harvest by allowing only a specific number of individuals the opportunity to hunt.

The illegal kill of black bears documented by this study occurred exclusively in females; however, some of the unknown or suspect mortality in both sexes may have also been illegal kills. Illegal kills in this study represented not only the loss of the adult female, but also the loss of their newborn cubs. Under Montana law, it is currently legal to carry a loaded weapon in a vehicle while hunting. Shooting from a public highway is prohibited, but U.S. Forest Service roads are not considered public highways. Changes in these laws and an intensive hunter information and education program could reduce the mortality of females with cubs, young bears, and wounding losses often associated with quick shots. Increased law enforcement activity through additional personnel, patrols, undercover operations, rewards for information, and vigorous prosecution could decrease illegal take through deterrent effects.



LITERATURE CITED

- Alt, G. 1982. Reproductive biology of Pennsylvania's black bears. Pa. Game News 53(2):9-15.
- Aune, K. and W. Kasworm. 1989. East Front grizzly bear study; final report. Mont. Dept. Fish, Wildl. & Parks. Helena. 332pp.
- Beecham, J. 1980. Population characteristics, denning, and growth patterns of black bears in Idaho. PhD. Diss., Univ. Montana, Missoula. 110pp.
- Brown, G. W., R. B. Campbell, H. J. Cross, T. O. Lemke, and S. J. Riley. 1987. P-R Job Prog. Rept. W-130-R-18, I-1. Montana Dept. Fish, Wildl. & Parks, Helena. 83pp.
- Bunnell, F. L., and D. E. N. Tait. 1980. Bears in models and in reality- implications to management. Int. Conf. Bear Res. and Manage. 4:15-24.
- Bunnell, F. L., and D. E. N. Tait. 1985. Mortality rates of North American bears. Arctic 38:318-323.
- Downing, R. L. 1980. Vital statistics of animal populations. Pages 247-267 in Wildlife Management Techniques Manual. S. D. Schemnitz, ed. The Wildlife Society. Washington, D.C. 686pp.
- Heisey, D. M. and T. K. Fuller. 1985. Evaluation of survival and cause-specific mortality rates using telemetry data. J. Wildl. Manage. 49:668-674.
- Hellgren, E. C., D. W. Carney, N. P. Garner, and M. R. Vaughan. 1988. Use of breakaway cotton splicers on radio collars. Wildl. Soc. Bull. 16:216-218.
- Hugie, R. D. 1982. Black bear ecology and management in the northern conifer deciduous forests of Maine. PhD diss., Univ. of Montana, Missoula. 217pp.
- Johnson, K. G. and M. R. Pelton. 1980. Prebaiting and snaring techniques for black bears. Wildl. Soc. Bull. 8:46-54.
- Jonkel, C. J. and I. McT. Cowan. 1971. The black bear in the spruce-fir forest. Wildl. Monogr. 27. 57pp.
- Kasworm, W. and T. Manley. 1988. Cabinet Mountains grizzly bear study. Montana Dept. Fish, Wildl. & Parks, Helena. 81pp.
- \_\_\_\_\_. and \_\_\_\_\_. 1990. Road and trail influences on grizzly bears and black bears in northwest Montana. Int. Conf. Bear Res. and Manage. 8:79-84.
- McLellan, B. N. 1988. Dynamics of a grizzly bear population during a period of industrial resource extraction. III. Natality and rate of increase. Canadian J. Zool. 67:1865-1868.

Roge

Ston

Stri

Wadd



- Rogers, L. L. 1987. Effects of food supply and kinship on social behavior, movements, and population growth of black bears in northeastern Minnesota. Wildl. Monogr. 97. 72pp.
- Stoneberg, R. and C. J. Jonkel. 1966. Age determination in black bears by cementum layers. J. Wildl. Manage. 30:411-414.
- Stringham, S. F. 1990. Black bear reproductive rate relative to body weight in hunted populations. Int. Conf. Bear Res. and Manage. 8:425-432.
- Waddel, T. E. and D. E. Brown. 1984. Exploitation of two subpopulations of black bears in an isolated mountain range. J. Wildl. Manage. 48:933-938.



Appendix Table 1. Capture information, monitoring periods, and fate of radio collared black bears in northwest Montana, 1983-1990.

Tag Number	Sex	Capture Age	District	Capture Date	Days Monitored	Date of Last Relocation	Fate
84-85	F	4	104	6 May 83	728	3 May 85	Suspect Mortality
88-89	M	5	104	11 May 83	1,524	13 Jul 87	Radio Failure
90-91	M	12	104	13 May 83	702 <sup>1</sup>	14 Apr 85	Radio Failure
263-264	M	10	104	23 May 83	342	29 Apr 84	Hunter Kill
366-367	M	10	121	29 May 83	716	14 May 85	Hunter Kill
686-687	F	15	121	26 May 84	871	13 Oct 86	Mortality (Unknown Cause)
701-702	F	21	121	28 May 84	1,136	7 Jul 87	Natural Mortality
726-727	F	6	121	29 May 84	1,781	14 Apr 89	Alive
736-737	F	7	121	18 Jun 84	1,031	14 Apr 87	Radio Failure
763-764	M	16	104	24 Jun 84	320	9 May 86	Hunter Kill
799-800	F	9	104	28 Jun 84	656 <sup>1</sup>	14 Apr 86	Radio Failure
779-780	F	11	121	11 Oct 84	1,150	14 Apr 89	Alive
630-631	M	16	104	10 May 85	97	14 Aug 85	Lost Collar
757-758	F	13	104	16 May 85	463	21 Aug 86	Suspect Mortality
11-11	F	8	104	16 May 85	421	10 Jul 87	Suspect Mortality
13-13	F	10	104	24 May 85	362	20 May 86	Illegal Kill (with cubs)
7-7	F	7	104	6 Jun 85	689	25 Apr 87	Hunter Kill
265-266	M	12	104	7 Jun 85	701	8 May 87	Hunter Kill
5-5	F	7	104	15 Jun 85	1,034	14 Apr 85	Radio Failure
10-10	F	14	104	22 Jun 85	343	30 May 86	Illegal Kill (lactating)
26-26	F	11	104	12 Jul 85	308	15 May 86	Hunter Kill
102-102	M	8	100	4 May 86	1,144	21 Jun 89	Lost Collar
103-103	M	7	100	10 May 86	367	11 May 87	Hunter Kill
104-104	M	8	100	12 May 86	357	3 May 87	Hunter Kill
31-31	M	11	104	12 May 86	427	12 Jul 87	Lost Collar
105-105	F	5	100	22 May 86	1,058	14 Apr 89	Radio Failure
110-110	F	15	100	30 May 86	141	17 Oct 86	Mortality (Unknown cause)
111-111	M	9	100	5 Jun 86	342	12 May 87	Hunter Kill
112-112	M	7	100	13 Jun 86	389	6 Jul 87	Lost Collar
37-37	M	11	121	15 Jun 86	354	3 Jun 87	Suspect Mortality
255-256	F	9	104	3 Jul 86	1,016	14 Apr 89	Alive
42-42	F	9	104	9 Jul 86	1,010	14 Apr 89	Alive
41-41	F	10	104	11 Jul 86	1,008	14 Apr 89	Alive
120-120	F	8	100	30 Aug 86	1,160	2 Nov 89	Mortality (Unknown cause)
123-123	F	7	100	12 Sep 86	240	9 May 87	Hunter Kill
130-130	M	4	100	26 May 87	1,284	30 Nov 90	Alive
132-132	M	6	100	9 Jun 87	703	13 May 89	Hunter Kill
133-133	F	7	100	13 Jun 87	1,266	30 Nov 90	Alive
114-135	M	6	100	1 Jul 87	409	13 Aug 88	Lost Collar
748-749	M	6	104	28 Aug 87	595 <sup>1</sup>	14 Apr 89	Alive
186-186	F	8	100	30 Sep 89	427	30 Nov 90	Alive
191-191	F	4	100	12 Oct 89	202	1 May 90	Hunter Kill
117-198	M	9	100	30 May 90	185	30 Nov 90	Alive
219-219	F	15	100	19 Sep 90	73	30 Nov 90	Alive
220-220	F	5	100	25 Sep 90	67	30 Nov 90	Alive

<sup>1</sup>Collar used previously on another animal.



Appendix Table 2. Age and sex specific life tables for captured male and female black bears in hunting districts 100, 104, and 121, 1983-1990.

-----  
Males  
-----

Age	Hunting District 100				Hunting District 103				Hunting District 121			
	Number Caught	Initial Cohort	Lx	Survival Rate	Number Caught	Initial Cohort	Lx	Survival Rate	Number Caught	Initial Cohort	Lx	Survival Rate
4	6	39	1.000	0.846	11	54	1.000	0.796	3	27	1.000	0.889
5	4	33	0.846	0.879	5	43	0.796	0.884	3	24	0.889	0.875
6	5	29	0.744	0.828	8	38	0.704	0.789	5	21	0.778	0.762
7	4	24	0.615	0.833	7	30	0.556	0.767	5	16	0.593	0.688
8	4	20	0.513	0.800	3	23	0.426	0.870	2	11	0.407	0.818
9	2	16	0.410	0.875	7	20	0.370	0.650	2	9	0.333	0.771
10	3	14	0.359	0.786	2	13	0.241	0.846	3	7	0.259	0.571
11	3	11	0.282	0.727	3	11	0.204	0.727	0	4	0.148	1.000
12	0	8	0.205	1.000	4	8	0.148	0.500	0	4	0.148	1.000
13	2	8	0.205	0.750	1	4	0.074	0.750	0	4	0.148	1.000
14	4	6	0.154	0.333	0	3	0.056	1.000	0	4	0.148	1.000
15	1	2	0.051	0.500	1	3	0.056	0.667	1	4	0.148	0.750
16	0	1	0.026	1.000	1	2	0.037	0.500	1	3	0.111	0.667
17	0	1	0.026	1.000	1	1	0.019	0.000	0	2	0.074	1.000
18	1	1	0.026	0.000					0	2	0.074	1.000
19									1	2	0.074	0.500
20									0	1	0.037	1.000
21									0	1	0.037	1.000
22									1	1	0.037	0.000
Mean Survival Rate			0.744				0.696				0.805	

-----  
Females  
-----

Age	Hunting District 100				Hunting District 103				Hunting District 121			
	Number Caught	Initial Cohort	Lx	Survival Rate	Number Caught	Initial Cohort	Lx	Survival Rate	Number Caught	Initial Cohort	Lx	Survival Rate
4	2	21	1.000	0.905	4	36	1.000	0.889	1	12	1.000	0.917
5	2	19	0.905	0.895	5	32	0.889	0.844	0	11	0.917	1.000
6	1	17	0.810	0.941	3	27	0.750	0.889	3	11	0.917	0.727
7	4	16	0.762	0.750	5	24	0.667	0.792	1	8	0.667	0.875
8	6	12	0.571	0.500	3	19	0.528	0.842	0	7	0.583	1.000
9	2	6	0.286	0.667	3	16	0.444	0.813	1	7	0.583	0.857
10	1	4	0.190	0.750	2	13	0.361	0.846	0	6	0.500	1.000
11	1	3	0.143	0.667	3	11	0.306	0.727	1	6	0.500	0.833
12	0	2	0.095	1.000	4	8	0.222	0.500	2	5	0.417	0.600
13	0	2	0.095	1.000	0	4	0.111	1.000	0	3	0.250	1.000
14	0	2	0.095	1.000	3	4	0.111	0.250	0	3	0.250	1.000
15	2	2	0.095	0.000	0	1	0.028	1.000	2	3	0.250	0.333
16					1	1	0.028	0.000	0	1	0.083	1.000
17									0	1	0.083	1.000
18									0	1	0.083	1.000
19									0	1	0.083	1.000
20									0	1	0.083	1.000
21									0	1	0.083	1.000
									1	1	0.083	0.000
Mean Survival Rate			0.756				0.722				0.841	



Appendix Table 3. Age and sex specific life tables for hunter harvested male and female black bears in hunting districts 100, 104, and 121, 1985-1989.

Males

Age	Hunting District 100				Hunting District 103				Hunting District 121			
	Number Caught	Initial Cohort	Lx	Survival Rate	Number Caught	Initial Cohort	Lx	Survival Rate	Number Caught	Initial Cohort	Lx	Survival Rate
4	24	113	1.000	0.752	21	117	1.000	0.821	39	144	1.000	0.729
5	20	85	0.752	0.765	14	86	0.821	0.854	14	105	0.729	0.867
6	11	65	0.575	0.831	13	82	0.701	0.841	22	91	0.632	0.758
7	10	54	0.478	0.815	9	69	0.590	0.870	12	69	0.479	0.826
8	7	44	0.389	0.841	11	60	0.513	0.817	10	57	0.396	0.825
9	7	37	0.327	0.811	11	49	0.419	0.776	8	47	0.326	0.830
10	7	30	0.265	0.767	4	38	0.325	0.895	8	39	0.271	0.795
11	11	23	0.204	0.522	7	34	0.291	0.794	8	31	0.215	0.742
12	2	12	0.106	0.833	2	27	0.231	0.926	7	23	0.160	0.696
13	1	10	0.088	0.900	6	25	0.214	0.760	1	16	0.111	0.938
14	2	9	0.080	0.778	9	19	0.162	0.526	2	15	0.104	0.867
15	4	7	0.062	0.429	2	10	0.085	0.800	4	13	0.090	0.692
16	2	3	0.027	0.333	2	8	0.068	0.750	1	9	0.063	0.889
17	0	1	0.009	1.000	1	6	0.051	0.833	1	8	0.056	0.875
18	1	1	0.009	0.000	1	5	0.043	0.800	0	7	0.049	1.000
19					1	4	0.034	0.750	2	7	0.049	0.714
20					1	3	0.026	0.667	2	5	0.035	0.600
21					1	2	0.017	0.500	1	3	0.021	0.667
22					1	1	0.009	0.000	0	2	0.014	1.000
23									1	2	0.014	0.500
24									1	1	0.007	0.000
Mean Survival Rate			0.692					0.736				0.753

Females

Age	Hunting District 100				Hunting District 103				Hunting District 121			
	Number Caught	Initial Cohort	Lx	Survival Rate	Number Caught	Initial Cohort	Lx	Survival Rate	Number Caught	Initial Cohort	Lx	Survival Rate
4	16	68	1.000	0.765	13	76	1.000	0.829	9	99	1.000	0.909
5	10	52	0.765	0.808	8	63	0.829	0.873	12	90	0.909	0.867
6	4	42	0.618	0.905	7	55	0.724	0.873	4	78	0.747	0.949
7	11	38	0.559	0.711	8	48	0.632	0.833	9	74	0.747	0.878
8	5	27	0.397	0.815	8	40	0.526	0.800	7	65	0.657	0.892
9	4	22	0.324	0.818	8	32	0.421	0.750	13	58	0.586	0.776
10	5	18	0.265	0.722	4	24	0.316	0.833	10	45	0.455	0.778
11	1	13	0.191	0.923	5	20	0.263	0.750	7	35	0.354	0.800
12	3	12	0.176	0.750	1	15	0.197	0.933	9	28	0.283	0.679
13	1	9	0.132	0.889	4	14	0.184	0.714	5	19	0.192	0.737
14	1	8	0.118	0.875	1	10	0.132	0.900	4	14	0.141	0.714
15	3	7	0.103	0.571	3	9	0.118	0.667	3	10	0.101	0.700
16	2	4	0.059	0.500	3	6	0.079	0.500	1	7	0.071	0.857
17	0	2	0.029	1.000	2	3	0.039	0.333	2	6	0.061	0.667
18	0	2	0.029	1.000	0	1	0.013	1.000	2	4	0.040	0.500
19	1	2	0.029	0.500	0	1	0.013	1.000	0	2	0.020	1.000
20	1	1	0.015	0.000	1	1	0.013	0.000	1	2	0.020	0.500
21								1.000	1	1	0.010	0.000
Mean Survival Rate			0.738					0.741				0.733

SUM  
MON

Tim  
Lib

Abs  
spr  
as  
tot  
Com  
hun  
and  
hun  
num  
Suc  
tha  
was  
kil  
2.5  
the  
rea  
tha  
the  
Onl  
per  
of  
fro  
lea

nor  
deg  
pop  
of  
sea  
28  
tra  
of

que  
HD  
Alt  
hel  
con  
oth  
use



SUMMARY OF QUESTIONNAIRE GIVEN TO SPRING BLACK BEAR HUNTERS IN NORTHWESTERN MONTANA

Timothy J. Thier, U.S. Fish and Wildlife Service, 475 Fish Hatchery Road, Libby, MT 59923

Abstract: In 1986 and 1987, questionnaires were distributed to as many of the spring black bear (*Ursus americanus*) hunters in Hunting District 100 (HD 100) as possible, with a special effort made to contact all successful hunters. A total of 125 questionnaires were returned, 61 of them from successful hunters. Completed questionnaires were obtained from 78% of all successful spring hunters in HD 100, as recorded by the Montana Department of Fish, Wildlife, and Park's (MDFWP) mandatory tooth return program for successful hunters. All hunters were asked a variety of questions that related to hunter techniques, number and types of bears observed, and the amount of hunter effort. Successful hunters were asked to complete an additional section with questions that pertained to the time of day the bear was killed, the distance the bear was killed from an open road, and the type of habitat in which the bear was killed. The mean number of bears observed was 4.2 for successful hunters and 2.5 for unsuccessful hunters. Thirty-four percent of the respondents claimed they declined shooting at a legal bear, with "small size" the most common reason given. Twenty-four hunters stated they shot at a minimum of 29 bears that were not killed, of which at least 2 were wounded. "Driving roads" was the most common hunting technique given, with walking closed roads second. Only 6 of the 61 bears reported killed were shot before noon. Fifty-two percent of the successful hunters reported they killed their bear within 300 m of an open road. Sixty-six percent of the bears killed were first observed from an open road. Clearcuts and/or roads were stated as the habitat where at least 70% of the bears were killed.

---

In 1986 and 1987, data was collected on black bears in HD 100 of extreme northwestern Montana in partial fulfillment of the requirements for an M.S. degree in Environmental Studies (Thier 1990). That study addressed the population characteristics of black bears in the area and examined the impacts of hunting, primarily during the spring hunting season. The spring hunting season in HD 100 typically extends from 15 April to 15 May. During the study 28 individual black bears were captured, marked and released, 13 with radio transmitting collars. The movements of instrumented bears and the mortality of both marked and unmarked bears was carefully monitored.

In an effort to better understand hunter technique, effort, and success, questionnaires were distributed to as many of the spring black bear hunters in HD 100 as possible, with a special effort made to contact successful hunters. Although information from Montana's annually conducted telephone survey was helpful, it failed to address many important points. The objectives in conducting this survey was to gather information that might complement the other research being conducted, and to gather baseline data that would be useful in future management.